

Improving Pediatric Rapid Response Team Performance Through Crew Resource Management Training of Team Leaders

Ashley Siems, MD, Alexander Cartron, BS, Anne Watson, MSc, PhD, BSN, RN, Robert McCarter Jr, ScD, Amanda Levin, MD

ABSTRACT

BACKGROUND: Rapid response teams (RRTs) improve the detection of and response to deteriorating patients. Professional hierarchies and the multidisciplinary nature of RRTs hinder team performance. This study assessed whether an intervention involving crew resource management training of team leaders could improve team performance.

METHODS: In situ observations of RRT activations were performed pre- and post-training intervention. Team performance and dynamics were measured by observed adherence to an ideal task list and by the Team Emergency Assessment Measure tool, respectively. Multiple quartile (median) and logistic regression models were developed to evaluate change in performance scores or completion of specific tasks.

RESULTS: Team leader and team introductions (40% to 90%, $P = .004$; 7% to 45%, $P = .03$), floor team presentations in Situation Background Assessment Recommendation format (20% to 65%, $P = .01$), and confirmation of the plan (7% to 70%, $P = .002$) improved after training in patients transferred to the ICU ($n = 35$). The Team Emergency Assessment Measure metric was improved in all 4 categories: leadership (2.5 to 3.5, $P < .001$), teamwork (2.7 to 3.7, $P < .001$), task management (2.9 to 3.8, $P < .001$), and global scores (6.0 to 9.0, $P < .001$) for teams caring for patients who required transfer to the ICU.

CONCLUSIONS: Targeted crew resource management training of the team leader resulted in improved team performance and dynamics for patients requiring transfer to the ICU. The intervention demonstrated that training the team leader improved behavior in RRT members who were not trained.

www.hospitalpediatrics.org

DOI:10.1542/hpeds.2016-0111

Copyright © 2017 by the American Academy of Pediatrics

Address correspondence to Ashley Siems, MD, Department of Critical Care Medicine, Children's National Health System, 111 Michigan Ave NW, Washington, DC 20010. E-mail: asiems@cnmc.org

HOSPITAL PEDIATRICS (ISSN Numbers: Print, 2154-1663; Online, 2154-1671).

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: This work was supported by awards UL1TR000075 and KL2TR000076 from the National Institutes of Health National Center for Advancing Translational Sciences. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the National Center for Advancing Translational Sciences or the National Institutes of Health. Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

Dr Siems assisted with designing the study, designed and executed the crew resource management intervention, assisted with data analysis, and drafted the initial manuscript; Mr Cartron designed the study's data collection instrument and database, collected data, assisted with data analysis, and critically reviewed and revised the manuscript; Dr Watson assisted with designing the study, supervised and assisted with design of the data collection instrument, and critically reviewed and revised the manuscript; Dr McCarter performed statistical analysis and critically reviewed and revised the manuscript; Dr Levin conceptualized and designed the study, supervised design of the crew resource management intervention, assisted with data analysis, and critically reviewed and revised the manuscript; and all authors approved the final manuscript as submitted.



Children's National Health System, Washington, DC

In 2006, the Institute for Healthcare Improvement published their “100,000 Lives Campaign” and included rapid response teams (RRTs) as a modality to improve patient safety.¹ The adoption of RRTs has prevented cardiac and respiratory arrest, improved survival after these events, reduced the need for ICU-level treatments soon after transfer, and lessened the time between deterioration and treatments.²⁻⁹ In spite of these successes, RRTs, like other complex medical teams, are challenged by difficulties with communication and collaborative problem-solving.¹⁰ Barriers to RRT performance include varying levels of confidence with one’s own skills, challenges overcoming professional hierarchies, and expectations of adverse interpersonal conflict.¹¹ Recently, the 2016 Joint Commission National Patient Safety Goal to “improve the effectiveness of communication among caregivers” signaled the growing recognition that improvement in the “nontechnical” social interaction skills of health care providers is critical for the delivery of safe and effective patient care.¹² Adverse events can often be attributed to poor interpersonal interactions and coordination rather than insufficient clinical knowledge.^{13,14} Communication errors have been reported to be responsible for nearly 60% of all medical errors.¹⁵ Thus, approaches to improving RRT performance are necessary and should focus on interpersonal interactions and communication.

One such approach is crew resource management (CRM), which is a strategy of training with a focus on nontechnical skills that complement a provider’s clinical knowledge and promote the delivery of safe care.¹⁶ CRM training of high-performance teams in other high-risk industries, including the military, aviation, and nuclear power fields, includes adoption of standardized tools and behaviors to improve teamwork and reduce risk.¹⁷⁻²⁴ CRM focuses on leadership, problem-solving, situational awareness, communication skills, and resource management to enhance team performance.²⁵⁻²⁷ Careful consideration of one’s own fatigue coupled with verbal and nonverbal communication are paramount. CRM training is most

effective when structured around simulation exercises, in which providers have the opportunity to safely practice vital nontechnical skills and receive real-time feedback in an environment uncoupled from adverse events.^{28,29} Although nontechnical skills have not historically been taught in clinical medical education, implementation of CRM training in the ICU and other medical settings has been shown to reduce complications and lower mortality in critically ill patients.^{30,31} Due to its previous successful application in high-risk industry and preliminary data suggesting it can be applied to medical teams, we hypothesized that CRM training can help improve RRT performance. The purpose of the current study was to evaluate CRM nontechnical skills training as an improvement technique for RRT performance.

In designing this study, we recognized that training and maintaining the skills of every potential responder to an RRT (nurses, trainees, rotating trainees, and respiratory therapists) was expensive and impractical to sustain. We targeted the team leaders of our RRT to have a high impact and minimize the cost and time barriers associated with comprehensive CRM training programs.^{32,33} We hypothesized that training the team leader would improve the overall team performance as well as completion of ideal tasks during RRT activation.

METHODS

Setting and Design

This study was conducted on the acute care floors of a single, urban, tertiary 313-bed pediatric medical center from March 2015 to December 2015. This institution averages 50 RRT activations per month. The RRT consists of personnel from critical care medicine (CCM) including a critical care fellow or nurse practitioner (NP), an ICU charge nurse, and an ICU respiratory therapist (RT) who joins the acute care floor team consisting of the bedside nurse, floor charge nurse, primary team physician or physician extender, and floor RT at the patient’s bedside. A pre- and postintervention design was used to assess leadership and team performance during RRT activations. Observations of RRT activations occurred before and after team

leader training in CRM. Observations were conducted in situ to assess the impact of our training in real-life hospital environments with challenges of stress, time-pressure, and coordination of personnel unfamiliar with one another.

Team Leader Training

Our team leader training consisted of 2 sessions that were conducted by physician champions with knowledge of CRM principles. The first 90-minute training session was built on adult learning theory.³⁴ The learners engaged in active learning and were introduced to evidence that rapid response systems reduce patient morbidity while being instructed on the ideal RRT task list and CRM principles. First-year CCM fellows and NPs attended the first session because it was designed to target junior practitioners. The second 60-minute session used simulation training to reinforce CRM principles and apply them to common RRT scenarios. All CCM fellows and NPs attended the second session. We defined junior practitioners as NPs and first-year fellows because these providers were new to both our ICU and to the team leader role on the RRT.

Observations

Members of the research team (A.C., A.W.) conducted all in situ observations based on the observer’s availability when the RRT system was activated. RRTs at our institution are activated via pagers. The observers carried pagers and would go directly to the patient room when an RRT was activated. The ICU team has 15 minutes to respond to the page allowing for the observer to be in place before the ICU team arrival. All observations were during the day and represent a convenience sampling.

The RRT observations were conducted in 3 phases. Ten RRTs were observed in the project development phase to validate and test the data collection instrument. These observations were not included in the final analysis. During this phase, both observers were present at each RRT observed. The second phase was the preintervention stage, where 36 RRT activations were observed during the spring and early summer of 2015. The intervention, team

leader training, occurred during the midsummer of 2015 with 37 postintervention observations occurring in the late summer and fall. During the second and third phase, only 1 observer was required to be present, although 20% of observations in these phases had both observers present. Interrater reliability was assessed using the observations that both researchers attended.

The observers used a paper data collection tool and recorded patient demographics and characteristics of the RRT events (eg, patient age, sex, mean duration of the event, and who called the RRT). The disposition status was also recorded to determine which patients were transferred to the ICU. The observers are not members of the RRT and although they may be identified by RRT members as researchers, the nature and purpose of their observations were not disclosed until the end of the study.

Measures of Leadership and Team Performance

Two metrics were used to assess team performance: 1 to observe the completion of tasks essential to RRT activations and a second to assess the nontechnical skills of the RRT. A focus group of key stakeholders in RRTs at our institution, including ICU physicians, nurses, RTs, acute care nurses, nurse educators, and acute care physicians, identified 12 ideal tasks essential to RRT function. The task list was developed from consensus opinion and previously published research. Elements previously shown to enhance team performance and patient outcomes, including using standard safety language such as Situation Background Assessment Recommendation (SBAR) to enhance shared mental models, formulating and stating a clear plan if the patient is going to remain on the acute care floor, and using cross-check techniques to identify concerns with the plan key elements of the task list.^{28,35–38} The 12 ideal tasks were: (1) measuring patient vital signs; (2) ICU team introduces themselves; (3) ICU team leads team introductions; (4) acute care team presents using SBAR; (5) team leader asks for additional input from the acute care team; (6) ICU team assesses patient; (7) ICU

team formulates a clear plan with disposition; the plan delineates a time frame for (8) interventions, (9) task delegation, and (10) an escalation plan given with expectations of outcome; (11) acute care team repeats back plan; and (12) the ICU team asks for final concerns and questions.

The nontechnical skills of the team were scored using the Team Emergency Assessment Measure (TEAM), a 12-item tool validated to measure the performance of medical emergency teams.^{39–41} The TEAM tool includes 12 statements and offers subscore composites in leadership, team work, and task management measured on a 0 to 4 Likert scale, ranging from “never/hardly ever” (0) to “about as often as not” (2) to “always/nearly always” (4). Overall global team nontechnical performance was measured using a Likert scale ranging from 0 to 10.

Adherence to each item on the RRT ideal task list was recorded during the event. Within 15 minutes after the event, the observer rated the nontechnical skills of the team using the TEAM tool. Study data were managed using REDCap (Research Electronic Data Capture, Vanderbilt

University, Nashville, TN) tools hosted at Children’s National Medical Center.⁴²

Data Analysis

Sample size calculation was powered to detect a 15% improvement in the global rating of the TEAM score, resulting in an $N = 35$ per observation group. Descriptive statistics are presented as percentages for dichotomous variables, averages, and SDs. Pre and postintervention comparisons were presented as adherence frequencies $\pm 95\%$ confidence intervals (CI) for elements of the ideal RRT task list and as medians with 95% CIs for the overall and component TEAM scores. P values $< .05$ were regarded as statistically significant. We performed a quantile (median) regression model based on quantile regression in Stata (Stata Corp, College Station, TX) for the analysis of scores. This regression enabled us to evaluate interaction effects of patient severity defined by the need for ICU transfer. The distributions of study outcomes was not normal (parametric) and could not be transformed to meet the parametric assumption required for linear regression modeling. Estimating the median provided a nonbiased estimate of the central tendency of the sample distribution and by inference

TABLE 1 Patient Demographics and Characteristics of RRT Activations Pre- and Post-CRM Training

Characteristic	Pre ($n = 36$)	Post ($n = 37$)	P
Average patient age, mo (mean \pm SD)	91 \pm 77	73 \pm 69	.30
Patient sex			.35
Male, n (%)	21 (58)	17 (46)	
Patient insurance type (%)			
Non-Medicaid	9 (25)	10 (27)	.84
Medicaid	25 (75)	27 (73)	
English speaking (%)	32 (89)	28 (76)	.12
Transferred to ICU (%)	15 (42)	20 (54)	.35
Time of call (%)			.48
Morning	22 (61)	19 (51)	
Afternoon	14 (39)	18 (49)	
Mean time to arrival, min (mean \pm SD)	9.5 \pm 4.16	10.0 \pm 4.08	.57
Mean duration of call, min (mean \pm SD)	11.0 \pm 4.26	12.8 \pm 7.93	.26
Who called RRT? (%)			
Primary provider	17 (47)	19 (51)	.94
Nurse	16 (44)	15 (41)	
Other	3 (8)	3 (8)	
Parent/family present, n (%)	25 (69)	28 (76)	.61

TABLE 2 Percentage Adherence to Ideal RRT Tasks Pre- and Post-CRM Training

	Remained on Floor % (95% CI) (n = 38)			Transferred to ICU % (95% CI) (n = 35)		
	Pre (n = 21)	Post (n = 17)	P	Pre (n = 15)	Post (n = 20)	P
PICU fellow/NP introduces self, % (95% CI)	62 (41–82)	88 (72–100)	.08	40 (15–65)	90 (77–100)	.004
PICU fellow/NP leads introduction of team, % (95% CI)	14 (0–29)	41 (17–64)	.07	7 (0–19)	45 (23–67)	.03
Primary RN or LIP presents SBAR, % (95% CI)	28 (9–48)	64 (42–87)	.03	20 (0–40)	65 (44–86)	.01
Robust plan stated (patient remained on floor), % (95% CI)	62 (41–83)	70 (48–92)	.57	N/A	N/A	N/A
Primary resident repeats the plan (patient remained on floor), % (95% CI)	38 (17–59)	53 (29–77)	.36	N/A	N/A	N/A
Cross-check, % (95% CI)	38 (17–59)	65 (42–87)	.11	7 (0–19)	70 (50–90)	.002
Update family, % (95% CI)	52 (31–74)	59 (35–82)	.69	67 (43–91)	80 (62–98)	.38

LIP is a licensed individual provider and may be a resident, NP, or attending physician.

of the population from which the sample was drawn. When we evaluated discreet (categorical) outcomes, we relied on logistic regression, which allowed us to estimate the change in odds of the outcome associated with a unit change in the independent variable (predictor) of interest. All statistics were performed on Stata 2013 software.

RESULTS

Interrater reliability was determined to be high in all categories, including adherence to ideal RRT tasks (intraclass correlation coefficient [ICC] = 0.97), TEAM rating in leadership (ICC = 0.98), team work (ICC = 0.99), task management (ICC = 0.96), and global rating (ICC = 0.97).

Patient demographics and characteristics of the RRT events are listed Table 1. Leader demographics pre- and postintervention were uniform with 56% of the RRT activations led by junior practitioners

preintervention and 51% postintervention ($P = .81$).

The regression model showed a significant interaction effect when grouping the events in terms of disposition: transfer versus no transfer to the ICU. Due to this interaction, adherence rates for items on the ideal task list pre- and postintervention are presented in Table 2 by transferred versus nontransferred. After the team leader training, the team leader introduced themselves (40% preintervention vs 90% postintervention, $P = .004$) and led introductions of the team more often (7% preintervention and 45% postintervention, $P = .03$) for patients who required ICU transfer as highlighted in Fig 1. Although no member of the floor team participated in CRM training or had been trained with the task list, floor team members presented their patient concerns using SBAR more often in both the ICU (20% preintervention to 65% postintervention, $P = .01$) and

nontransfer (28% preintervention to 64% postintervention, $P = .03$) groups postintervention. For patients who were transferred to the ICU, the occurrence of cross-check behavior improved 10-fold, from 7% preintervention to 70% postintervention ($P = .002$).

Evaluation of nontechnical skills and team dynamics using the TEAM tool is displayed in Table 3. All 3 composite measures (leadership, 2.5 [95% CI, 2.1–2.9] to 3.5 [95% CI, 3.1–3.9]; teamwork, 2.7 [95% CI, 2.3–3.1] to 3.7 [95% CI, 3.4–4.1]; task completion, 2.9 [95% CI, 2.5–3.2] to 3.8 [95% CI, 3.5–4.1]) and the global rating (6.0 [95% CI, 4.9–7.1] to 9.0 [95% CI, 8.0–10.0]) increased for the events with patient transfer to the ICU. In addition, the task management composite score for the events with no transfer to the ICU (3.3 [95% CI, 3.0–3.6] to 3.8 [95% CI, 3.5–4.2]) improved significantly after the CRM intervention. The results for the events with an ICU transfer are highlighted in Fig 2.




FIGURE 1 Improvement in RRT ideal tasks adhered to pre- and post-CRM training in patients transferred to the ICU. Error bars represent 95% CIs.

TABLE 3 Median TEAM Composite Subscores Pre- and Post-CRM Training

	Remained on Floor (<i>n</i> = 38)			Transferred to ICU (<i>n</i> = 35)		
	Pre (<i>n</i> = 21)	Post (<i>n</i> = 17)	<i>P</i>	Pre (<i>n</i> = 15)	Post (<i>n</i> = 20)	<i>P</i>
Leadership composite, median (95% CI)	3.5 (3.2–3.8)	3.5 (3.1–3.9)	1.0	2.5 (2.1–2.9)	3.5 (3.1–3.9)	<.001
Teamwork composite, median (95% CI)	3.4 (3.1–3.8)	3.7 (3.3–4.1)	.28	2.7 (2.3–3.1)	3.7 (3.4–4.1)	<.001
Task management composite, median, (95% CI)	3.3 (3.0–3.6)	3.8 (3.5–4.2)	.03	2.9 (2.5–3.2)	3.8 (3.5–4.1)	<.001
Global rating, median (95% CI)	8 (7.0–9.0)	8 (6.9–9.1)	1.0	6 (4.9–7.1)	9 (8.0–10.0)	<.001

DISCUSSION

In this pre- and postintervention direct observational study of team dynamics, we implemented a short, pragmatic, and cost-efficient CRM training session for the team leaders. We were able to demonstrate improved leadership and overall team performance of our RRT in the real hospital environment with the use of skills learned at the simulation center to crises in real clinical settings.⁴³

Significant effects of this training were found primarily for RRTs caring for patients who were transferred to the ICU, with poorer performance scores before the intervention compared with events where the patient was not transferred. In other words, postintervention scores in patients transferred to the ICU reached levels similar to the preintervention, non-ICU transfer group. Higher acuity patients may increase the stress of a situation, create more chaos, and lead to team breakdown.^{44,45} We believe that focusing education on nontechnical skills allowed team leaders to gain insight into the impact of their tone and leadership style during events, especially when the

outcome was transfer to the ICU. Concrete steps, such as leading introductions and facilitating an SBAR presentation, may have diffused the stress and chaos of a team trying to care for a rapidly deteriorating patient. The leaders may have also learned how to create a safe environment and reduce the burden of hierarchies on team performance.

Interestingly, we would have postulated that RRTs caring for patients who remain on the floor after RRT involvement might have had poorer team performance based on previous research.^{46–50} ICU team members often perceive that floor patients do not require ICU expertise, transfer occurs primarily due to nurse staffing issues on the acute care floor, and leaving the ICU for a RRT call compromises the care of patients in the ICU. Floor team members often perceive that they have difficulty accessing the resources they need on the floor, have received punitive responses from the ICU team for activating the RRT, and experience stress involved in having to justify a transfer. More research that specifically examines the flow of the 2 types of events is

needed to additionally refine nontechnical skill development interventions.

RRTs are fast-start, multidisciplinary teams composed of members from different professions and specialties who need to quickly establish trust and good communication in high pressure situations. Most implementations of CRM training occur in small fixed teams and include all team members. We adapted CRM to unfixed teams and trained only 1 vital team member, the leader. Although full-staff training in CRM principles has clear benefits for patient safety, the start-up costs balanced by concerns over efficacy make comprehensive implementation unfeasible at most institutions.^{31–33,51–53} We were able to incorporate a reproducible, short, and sustainable training program into an existing yearly critical care medicine education structure. This also helps to address concerns for training new staff and accounts for turnover. Finally, this strategy helps to reinforce the training before the skills expire.

By training the team leader in CRM principles, our data suggest that the entire




FIGURE 2 Bar and whisker plot for median TEAM composite subscores pre- and post-CRM training in patients transferred to the ICU.

team performance improved. Targeting a high impact person and affecting team behavior have also been studied in cardiac arrest teams with similar results.³² The task management composite subscore of the TEAM instrument was higher posttraining for both the patients who were transferred and those who were not. In addition, one of the ideal tasks, presentation of the event in SBAR format, improved after training of the leader without any training of the frontline providers for both groups. It may be that the leader provided prompts for this type of complete report, especially the “recommendation” component of SBAR, which may be difficult in hierarchical environments, such as a standard RRT activation. This finding suggests that training communication skills, including how to create a shared mental model, resulted in a safer environment for nurses and trainees to speak up and express their perspective.

There are certain limitations to this study. The Hawthorne effect, the potential change in behavior of the team leader due to awareness that their behavior was being observed, especially after the intervention, was a limitation that we anticipated. We aimed to minimize this by blinding RRT members to the aims of the study and specific metrics used in observation. Furthermore, it is common for our RRT team leaders to have experience with a varied number of team members with other types of observers, such as trainees, new staff, or administrative personnel, depending on the time of day. Another limitation we anticipated was the lack of independence of the ideal task list and TEAM metric, because both were assessed by the same observer. If the team completed the expected tasks on the list, it is plausible that the observers rated the TEAM metric higher. We attempted to limit this bias by using a strict data dictionary a priori for the TEAM metric Likert scores. Finally, the observations were conducted during normal business hours as part of a convenience sample due to research staff availability. Future research that observes RRT activations at night, when fatigue and decreased provider availability are likely, may give additional insight into team dynamics and performance.

Ultimately, we observed an improvement in objective task completion as well as in a validated metric for team performance. Our findings support the evidence that CRM improves nontechnical skills and add that CRM can be applied to RRTs. These findings help highlight the benefit of nontechnical skill education in the trainee curriculum.

CONCLUSIONS

Two short, successive simulation-based CRM trainings of RRT team leaders effectively improved team dynamics during RRTs for patients who were transferred to the ICU. The training also improved completion of ideal tasks during RRT activations and affected the behavior of those trained and untrained. Targeting high impact team members and training them in CRM principles improved the performance of the entire team.

Acknowledgments

We thank Abdullah Al Shammari, who worked on the manuscript, provided valuable background research, and contributed to the creation of the observation tool.

REFERENCES

- Berwick DM, Calkins DR, McCannon CJ, Hackbarth AD. The 100,000 lives campaign: setting a goal and a deadline for improving health care quality. *JAMA*. 2006;295(3):324–327
- Hayes LW, Dobyns EL, DiGiovine B, et al. A multicenter collaborative approach to reducing pediatric codes outside the ICU [published correction appears in *Pediatrics*. 2012;130(1):168–169]. *Pediatrics*. 2012;129(3). Available at: www.pediatrics.org/cgi/content/full/129/3/e785
- Chan PS, Jain R, Nallmothu BK, Berg RA, Sasson C. Rapid response teams: a systematic review and meta-analysis. *Arch Intern Med*. 2010;170(1):18–26
- Bonafide CP, Localio AR, Song L, et al. Cost-benefit analysis of a medical emergency team in a children's hospital. *Pediatrics*. 2014;134(2):235–241

- Bonafide CP, Localio AR, Roberts KE, Nadkarni VM, Weirich CM, Keren R. Impact of rapid response system implementation on critical deterioration events in children. *JAMA Pediatr*. 2014;168(1):25–33
- Hunt EA, Zimmer KP, Rinke ML, et al. Transition from a traditional code team to a medical emergency team and categorization of cardiopulmonary arrests in a children's center. *Arch Pediatr Adolesc Med*. 2008;162(2):117–122
- Hanson CC, Randolph GD, Erickson JA, et al. A reduction in cardiac arrests and duration of clinical instability after implementation of a paediatric rapid response system. *Postgrad Med J*. 2010;86(1015):314–318
- Famolare N, Romano JC. The implementation of PEARS training: supporting nurses in non-critical care settings to improve patient outcomes. *J Pediatr Nurs*. 2013;28(3):267–274
- Theilen U, Leonard P, Jones P, et al. Regular in situ simulation training of paediatric medical emergency team improves hospital response to deteriorating patients. *Resuscitation*. 2013;84(2):218–222
- Kitto S, Marshall SD, McMillan SE, et al. Rapid response systems and collective (in)competence: an exploratory analysis of intraprofessional and interprofessional activation factors. *J Interprof Care*. 2015;29(4):340–346
- Roberts KE, Bonafide CP, Paine CW, et al. Barriers to calling for urgent assistance despite a comprehensive pediatric rapid response system. *Am J Crit Care*. 2014;23(3):223–229
- The Joint Commission. National Patient Safety Goals Effective January 1, 2016. Available at: www.jointcommission.org/assets/1/6/2016_NPSG_HAP.pdf.
- Kohn LT, Corrigan JM, Molla S, eds. *To Err Is Human: Building a Safer Health System*. Washington, DC: The National Academies Press; 2000

14. Makary MA, Daniel M. Medical error—the third leading cause of death in the US. *BMJ*. 2016;353(2139)
15. Gong Y, Zhu M, Li J, Turley JP, Zhang J. Clinical communication ontology for medical errors. *Stud Health Technol Inform*. 2007;129(pt 2):1007–1011
16. Gordon M, Darbyshire D, Baker P. Non-technical skills training to enhance patient safety: a systematic review. *Med Educ*. 2012;46(11):1042–1054
17. Leonard M, Graham S, Bonacum D. The human factor: the critical importance of effective teamwork and communication in providing safe care. *Qual Saf Health Care*. 2004;13(suppl 1):i85–i90
18. Mayer CM, Cluff L, Lin W-T, et al. Evaluating efforts to optimize TeamSTEPPS implementation in surgical and pediatric intensive care units. *Jt Comm J Qual Patient Saf*. 2011;37(8):365–374
19. Malec JF, Torsher LC, Dunn WF, et al. The mayo high performance teamwork scale: reliability and validity for evaluating key crew resource management skills. *Simul Healthc*. 2007;2(1):4–10
20. Hughes KM, Benenson RS, Krichthen AE, Clancy KD, Ryan JP, Hammond C. A crew resource management program tailored to trauma resuscitation improves team behavior and communication. *J Am Coll Surg*. 2014;219(3):545–551
21. Buljac-Samardzic M, Dekker-van Doorn CM, van Wijngaarden JDH, van Wijk KP. Interventions to improve team effectiveness: a systematic review. *Health Policy*. 2010;94(3):183–195
22. Fargen KM, Velat GJ, Lawson MF, Firment CS, Mocco J, Hoh BL. Enhanced staff communication and reduced near-miss errors with a neurointerventional procedural checklist. *J Neurointerv Surg*. 2013;5(5):497–500
23. Su L, Spaeder MC, Jones MB, et al. Implementation of an extracorporeal cardiopulmonary resuscitation simulation program reduces extracorporeal cardiopulmonary resuscitation times in real patients. *Pediatr Crit Care Med*. 2014;15(9):856–860
24. Kelleher DC, Carter EA, Waterhouse LJ, Parsons SE, Fritzeen JL, Burd RS. Effect of a checklist on advanced trauma life support task performance during pediatric trauma resuscitation. *Acad Emerg Med*. 2014;21(10):1129–1134
25. Helmreich RL. On error management: lessons from aviation. *BMJ*. 2000;320(7237):781–785
26. Flin R, Maran N. Identifying and training non-technical skills for teams in acute medicine. *Qual Saf Health Care*. 2004;13(suppl 1):i80–i84
27. Cheng A, Donoghue A, Gilfoyle E, Eppich W. Simulation-based crisis resource management training for pediatric critical care medicine: a review for instructors. *Pediatr Crit Care Med*. 2012;13(2):197–203
28. Hicks CM, Kiss A, Bandiera GW, Denny CJ. Crisis Resources for Emergency Workers (CREW II): results of a pilot study and simulation-based crisis resource management course for emergency medicine residents. *GJEM*. 2012;14(6):354–362
29. Fung L, Boet S, Bould MD, et al. Impact of crisis resource management simulation-based training for interprofessional and interdisciplinary teams: A systematic review. *J Interprof Care*. 2015;29(5):433–444
30. Blackwood J, Duff JP, Nettel-Aguirre A, Djođovic D, Joynt C. Does teaching crisis resource management skills improve resuscitation performance in pediatric residents? *Pediatr Crit Care Med*. 2014;15(4):e168–e174
31. Haerkens MHTM, Kox M, Lemson J, Houterman S, van der Hoeven JG, Pickkers P. Crew Resource Management in the Intensive Care Unit: a prospective 3-year cohort study. *Acta Anaesthesiol Scand*. 2015;59(10):1319–1329
32. Fernandez Castelao E, Boos M, Ringer C, Eich C, Russo SG. Effect of CRM team leader training on team performance and leadership behavior in simulated cardiac arrest scenarios: a prospective, randomized, controlled study. *BMC Med Educ*. 2015;15(1):116
33. Moffatt-Bruce SD, Hefner JL, Mekhjian H, et al. What is the return on investment for implementation of a Crew Resource Management program at an academic medical center [published online ahead of print September 29, 2015]? *Am J Med Qual*. doi:10.1177/1062860615608938
34. Ambrose SA, Bridges MW, DiPietro M, Lovett MC, Norman MK. *How Learning Works: Seven Research-Based Principles for Smart Teaching*. Hoboken, NJ: John Wiley & Sons; 2010.
35. Compton J, Copeland K, Flanders S, et al. Implementing SBAR across a large multihospital health system. *Jt Comm J Qual Patient Saf*. 2012;38(6):261–268
36. De Meester K, Verspuy M, Monsieurs KG, Van Bogaert P. SBAR improves nurse-physician communication and reduces unexpected death: a pre and post intervention study. *Resuscitation*. 2013;84(9):1192–1196
37. Haig KM, Sutton S, Whittington J. SBAR: a shared mental model for improving communication between clinicians. *Jt Comm J Qual Patient Saf*. 2006;32(3):167–175
38. Brady PW, Muething S, Kotagal U, et al. Improving situation awareness to reduce unrecognized clinical deterioration and serious safety events. *Pediatrics*. 2013;131(1). Available at: www.pediatrics.org/cgi/content/full/131/1/e298
39. Cooper S, Cant R, Porter J, et al. Rating medical emergency teamwork performance: development of the Team Emergency Assessment Measure (TEAM). *Resuscitation*. 2010;81(4):446–452
40. Rovamo L, Nurmi E, Mattila MM, Suominen P, Silvennoinen M. Effect of a simulation-based workshop on multidisciplinary teamwork of newborn emergencies: an intervention study. *BMC Res Notes*. 2015;8:671

41. Cooper S, Cant R, Connell C, et al. Measuring teamwork performance: Validity testing of the Team Emergency Assessment Measure (TEAM) with clinical resuscitation teams. *Resuscitation*. 2016;101: 97–101
42. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377–381
43. Boet S, Bould MD, Fung L, et al. Transfer of learning and patient outcome in simulated crisis resource management: a systematic review. *Can J Anaesth*. 2014;61(6):571–582
44. Pfaff MS. Negative affect reduces team awareness: the effects of mood and stress on computer-mediated team communication. *Hum Factors*. 2012; 54(4):560–571
45. Riskin A, Erez A, Foulk TA, et al. The impact of rudeness on medical team performance: a randomized trial. *Pediatrics*. 2015;136(3):487–495
46. Levin AB, Brady P, Duncan HP, Davis AB. Pediatric rapid response systems: identification and treatment of deteriorating children. *Curr Treat Options Pediatr*. 2015;1(1):76–89
47. Azzopardi P, Kinney S, Moulden A, Tibballs J. Attitudes and barriers to a Medical Emergency Team system at a tertiary paediatric hospital. *Resuscitation*. 2011;82(2):167–174
48. Williams DJ, Newman A, Jones C, Woodard B. Nurses' perceptions of how rapid response teams affect the nurse, team, and system. *J Nurs Care Qual*. 2011;26(3):265–272
49. Braaten JS. CE: Original research: hospital system barriers to rapid response team activation: a cognitive work analysis. *Am J Nurs*. 2015;115(2): 22–32; test 33; 47
50. Mackintosh N, Humphrey C, Sandall J. The habitus of 'rescue' and its significance for implementation of rapid response systems in acute health care. *Soc Sci Med*. 2014;120:233–242
51. Navarro-Martínez J, Gálvez C, Rivera-Cogollo MJ, Galiana-Ivars M, Bolufer S, Martínez-Adsuar F. Intraoperative crisis resource management during a non-intubated video-assisted thoracoscopic surgery. *Ann Transl Med*. 2015;3(8):111
52. Chen JG, Kemper AR, Odetola F, Cheifetz IM, Turner DA. Prevalence, characteristics, and opinions of pediatric rapid response teams in the United States. *Hosp Pediatr*. 2012;2(3): 133–140
53. Kemper PF, de Bruijne M, van Dyck C, Wagner C. Effectiveness of classroom based crew resource management training in the intensive care unit: study design of a controlled trial. *BMC Health Serv Res*. 2011;11(1):304

**Improving Pediatric Rapid Response Team Performance Through Crew
Resource Management Training of Team Leaders**

Ashley Siems, Alexander Cartron, Anne Watson, Robert McCarter, Jr and Amanda
Levin

Hospital Pediatrics 2017;7;88

DOI: 10.1542/hpeds.2016-0111 originally published online January 24, 2017;

Updated Information & Services	including high resolution figures, can be found at: http://hosppeds.aappublications.org/content/7/2/88
Supplementary Material	Supplementary material can be found at: http://hosppeds.aappublications.org/content/suppl/2017/01/17/7.2.88.DCSupplemental
References	This article cites 46 articles, 9 of which you can access for free at: http://hosppeds.aappublications.org/content/7/2/88.full#ref-list-1
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: https://shop.aap.org/licensing-permissions/
Reprints	Information about ordering reprints can be found online: http://classic.hosppeds.aappublications.org/content/reprints



**Improving Pediatric Rapid Response Team Performance Through Crew
Resource Management Training of Team Leaders**

Ashley Siems, Alexander Cartron, Anne Watson, Robert McCarter, Jr and Amanda
Levin

Hospital Pediatrics 2017;7;88

DOI: 10.1542/hpeds.2016-0111 originally published online January 24, 2017;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://hosppeds.aappublications.org/content/7/2/88>

Hospital Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 2012. Hospital Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 2154-1663.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

