

BRIEF REPORT

Multicenter Observational Study of the Use of Nebulized Hypertonic Saline to Treat Children Hospitalized for Bronchiolitis From 2008 to 2014

Joshua Davis, MD,^a Amy D. Thompson, MD,^a Jonathan M. Mansbach, MD, MPH,^b Pedro A. Piedra, MD,^c Kohei Hasegawa, MD, MPH,^d Ashley F. Sullivan, MS, MPH,^d Janice A. Espinola, MPH,^d Carlos A. Camargo Jr, MD, DrPH^d

ABSTRACT

OBJECTIVES: Among children hospitalized for bronchiolitis, we examined temporal trends in the use of hypertonic saline (HTS) and the characteristics associated with receiving this treatment.

METHODS: We conducted a secondary analysis of data from 2 large, multicenter prospective cohort studies that included young children hospitalized with bronchiolitis during 5 winter seasons (2008–2014). Our outcome was receipt of HTS any time during the preadmission visit or hospitalization. For comparison with the observed trends in HTS use, we conducted a PubMed literature review of studies evaluating HTS use for bronchiolitis. We classified publications according to their assessment of HTS efficacy (positive, negative, or neutral).

RESULTS: Among 2709 hospitalized children, 241 (8.9%) received HTS. There was marked variability in HTS use by site (0%–91%), with use more common among children admitted to the ICU than those treated on the ward (31% vs 15%). Over the study period, administration of HTS increased from 2% during the 2008–2009 season to 27% during the 2011–2012 season, but then it decreased to 11% during the 2013–2014 season. Before 2010, the number of PubMed HTS publications ranged from 0 to 3 articles per year, with all classified as either positive or neutral. The number of positive publications increased in 2010 ($n = 5$), whereas negative publications peaked in 2014 ($n = 4$).

CONCLUSIONS: Use of HTS in children hospitalized with bronchiolitis increased during the 2008 to 2012 winter seasons and then declined. These findings paralleled trends in the HTS literature, with positive articles encouraging HTS use in early years followed by a growing number of neutral and negative articles after 2012.



www.hospitalpediatrics.org

DOI: <https://doi.org/10.1542/hpeds.2017-0002>

Copyright © 2017 by the American Academy of Pediatrics

^aDepartment of Emergency Medicine, Nemours/Alfred I. duPont Hospital for Children, Wilmington, Delaware;
^bDepartment of Medicine, Boston Children's Hospital, and
^cDepartment of Emergency Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts; and
^dDepartments of Molecular Virology and Microbiology, and Pediatrics, Baylor College of Medicine, Houston, Texas

Address correspondence to Amy D. Thompson, MD, Emergency Department, Nemours/ Alfred I. duPont Hospital for Children, 1600 Rockland Rd, Wilmington, DE 19803. E-mail: amy.thompson@nemours.org

HOSPITAL PEDIATRICS (ISSN Numbers: Print, 2154-1663; Online, 2154-1671).

FINANCIAL DISCLOSURE: Dr Camargo received the grants mentioned below; the other authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: This study was supported by grants U01 AI-067693, U01 AI-087881, R01 AI-114552, and UG3 OD-023253 to Dr Camargo from the National Institutes of Health (Bethesda, MD). The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

Drs Davis and Thompson conceptualized and designed the current analysis, contributed to data collection, and drafted the manuscript; Drs Mansbach and Piedra contributed to the design of the original studies, contributed to data collection, and made critical revisions to the manuscript; Dr Hasegawa contributed to the design of the original studies and analysis of the results and made critical revisions to the manuscript; Ms Sullivan conceptualized and designed the original studies, supervised the data collection, and made critical revisions to the manuscript; Ms Espinola conceptualized and designed the current analysis, analyzed the results, and drafted the manuscript; Dr Camargo conceptualized and designed the original studies and current analysis, supervised both studies, and drafted the manuscript; and all authors approved the final manuscript for submission.

Bronchiolitis is the leading cause of hospitalization of young children in the United States.^{1,2} The mean cost of these hospitalizations is \$8500, with a median length-of-stay (LOS) of 2 to 3 days.^{3,4} The annual direct cost of bronchiolitis hospitalizations nationwide is estimated at \$543 million,⁴ with total charges in excess of \$1.6 billion.⁵ Therefore, reducing LOS in bronchiolitis could greatly reduce health care expenditure.

Currently, the mainstay of treatment is supportive, including supplemental oxygen, nasopharyngeal suctioning, intravenous fluids, and pressure-support ventilation, when necessary.⁵ Pharmacological treatment in bronchiolitis is highly variable.^{6–8} Several approaches have been tried, including the use of inhaled bronchodilators, inhaled and systemic corticosteroids, and ribavirin. The American Academy of Pediatrics (AAP) recommends against all of these approaches in the routine management of bronchiolitis.⁹ A more recently introduced treatment of severe bronchiolitis is the use of inhaled 3% or 5% hypertonic saline (HTS).^{10,11} Although several early, small studies supported HTS use in infants with bronchiolitis, the studies differed in their inclusion criteria, concentration of HTS, and concomitant bronchodilator use.^{12–14} In a 2013 Cochrane systematic review, HTS was shown to reduce inpatient LOS and clinical severity scores in both the inpatient and outpatient setting.¹⁰ However, this same review found no significant short-term change in bronchiolitis severity when HTS was given in the emergency department (ED).¹⁰ A recent randomized trial of hypertonic versus normal saline found no difference in LOS, severity scores, or need for supplemental oxygen among hospitalized infants with bronchiolitis.¹⁵ Although the most recent 2014 AAP guidelines recommend against HTS use in the ED, they state that HTS may be considered for infants hospitalized with bronchiolitis.⁹ The actual usage of HTS over the past decade is not known.

To address this knowledge gap, we analyzed multicenter, multiyear databases of children hospitalized with bronchiolitis to identify

temporal trends in the usage of this new treatment approach for severe bronchiolitis. We also examined the demographic and clinical characteristics associated with HTS administration. Although all patients were enrolled before the release of the 2014 AAP guidelines,⁹ we hypothesized that the use of HTS over time would parallel the results of published studies that either did or did not support this novel treatment. Therefore, we conducted a literature review on HTS use for bronchiolitis to compare with the temporal trends observed in our cohort studies.

METHODS

Study Design

We analyzed data from 2 large, prospective, multicenter studies of children hospitalized for bronchiolitis: the 30th Multicenter Airway Research Collaboration (MARC-30) and the 35th Multicenter Airway Research Collaboration (MARC-35)^{16,17} (Appendix Tables 1 and 2). A more detailed description has been published elsewhere.^{16,17} Both studies were observational in nature and data collection was coordinated by the Emergency Medicine Network in Boston, Massachusetts. The number of participating sites varied over 6 bronchiolitis seasons (November 1–April 30): 13 from 2007 to 2008 (MARC-30), 16 from 2008 to 2009 (MARC-30), 14 from 2009 to 2010 (MARC-30), 9 from 2011 to 2012 (MARC-35), 17 from 2012 to 2013 (MARC-35), and 6 from 2013 to 2014 (MARC-35).

Patient management was at the discretion of the on-site health care provider. All patients were enrolled within 24 hours of admission. Inclusion criteria were an attending physician's diagnosis of bronchiolitis, age <24 months (MARC-30) or <12 months (MARC-35), and parent or guardian informed consent. Exclusion criteria for both studies were previous enrollment, transfer to a participating hospital >48 hours after the original admission time, and declining the nasopharyngeal aspirate collection.^{16,17} Both studies were approved by all local institutional review boards. The details of patient data collection and viral testing have been previously described.^{16–21}

Literature Search

To compare the observed HTS usage to trends in the literature, we conducted a PubMed search in December 2014 using the keywords “bronchiolitis” and “hypertonic saline.” We included English language articles on HTS use in bronchiolitis in all clinical settings with no date limit. A single author (J.D.) then categorized the conclusions of each HTS article as positive, negative, neutral, or noninformative. A positive article was defined as one that showed clinical benefit (eg, decreased hospitalization rate, decreased LOS, improved clinical score, positive conclusion). Inversely, a negative article was defined as one reporting clinical harm or a negative conclusion. A neutral article was one in which HTS demonstrated neither clinical benefit nor harm. Lastly, noninformative articles were those that did not primarily discuss HTS use in hospitalized children with bronchiolitis.

Statistical Analyses

The primary outcome was HTS receipt at any point during the preadmission period or inpatient stay for bronchiolitis. Data for inpatient HTS treatment was unavailable during the 2007–2008 season of MARC-30; thus, the primary analysis was restricted to the 2008–2014 study period ($n = 2709$). Of these, 2179 (80%) children had their “preadmission” visit in the enrolling hospital's ED, whereas the remaining children were directly admitted from other locations (eg, another hospital, doctor's office). All analyses were performed by using Stata 14.1 (StataCorp, College Station, TX). Data are presented as proportions and medians with interquartile ranges (IQRs).

We examined trends in the use of HTS treatment by study season and compared them with trends in the bronchiolitis literature. Unadjusted associations between receipt of HTS treatment and patient characteristics were analyzed by using χ^2 , Fisher's exact test, and Wilcoxon rank sum test, as appropriate. A multivariable logistic regression was conducted to evaluate independent predictors of HTS treatment. The final regression model accounts for potential clustering by site, with

TABLE 1 Characteristics of Children Hospitalized for Bronchiolitis by Use of HTS from 2008 to 2014

Characteristics	All (N = 2709)		No HTS (N = 2468)		HTS Use (N = 241)		P
	n	%	n	%	n	%	
Enrollment season							<.001
2008–2009	839	31	821	33	18	7	
2009–2010	854	32	796	32	58	24	
2011–2012	148	5	108	4	40	17	
2012–2013	658	24	557	23	101	42	
2013–2014	210	8	186	8	24	10	
Site region							<.001
Northeast	519	19	439	18	80	33	
Midwest	386	14	375	15	11	5	
South	1182	44	1036	42	146	61	
West	622	23	618	25	4	2	
Age at enrollment (mo), median (IQR)	2709	3.7 (1.8–7.3)	2468	3.7 (1.7–7.4)	241	3.4 (1.8–6.0)	.18
Sex							.62
Male	1623	60	1475	60	148	61	
Female	1086	40	993	40	93	39	
Race and/or ethnicity							.04
Non-Hispanic white	1022	38	915	37	107	44	
Non-Hispanic black or African American	621	23	582	24	39	16	
Hispanic	961	36	875	36	86	36	
Other	99	4	90	4	9	4	
Insurance							<.001
Private	890	33	779	32	111	46	
Public	1712	64	1586	65	126	52	
None	88	3	84	3	4	2	
Weight (kg), median (IQR)	2694	6.2 (4.7–8.2)	2453	6.3 (4.7–8.2)	241	6.1 (4.7–7.9)	.54
Parental history of asthma	866	32	780	32	86	36	.20
Prenatal smoke exposure	401	15	371	15	30	13	.28
Premature birth (≤ 37 wk)	767	28	703	29	64	27	.50
ICU or special care at birth	609	23	551	22	58	24	.54
Postnatal smoke exposure (in same room)	249	9	232	9	17	7	.22
Ever attended daycare	561	21	500	20	61	25	.07
Previous overnight hospitalization	514	19	469	19	45	19	.90
History of breathing problems	593	22	544	22	49	20	.54
Breathing problem began <1 d before ED visit	130	5	113	5	17	7	.09
Child discomfort in 24 h before ED visit							.07
None/mild	498	18	464	19	34	14	
Moderate	1164	43	1065	43	99	41	
Severe	1042	39	935	38	107	45	
Took antibiotics for illness before ED visit	464	17	422	17	42	18	.89
Used inhaled bronchodilator for breathing problem before ED visit	1086	40	959	39	127	53	<.001
Used corticosteroids for breathing problem before ED visit	349	13	313	13	36	15	.32

TABLE 1 Continued

Characteristics	All (N = 2709)		No HTS (N = 2468)		HTS Use (N = 241)		P
	n	%	n	%	n	%	
Virology							
No. of infections detected							.12
0	128	5	123	5	5	2	
1	1782	66	1621	66	161	67	
≥2	799	29	724	29	75	31	
RSV	2042	75	1841	75	201	83	.002
Rhinovirus	660	24	606	25	54	22	.46
ED visit							
Initial heart rate (beats per minute), median (IQR)	2678	162 (148–176)	2441	162 (148–176)	237	164 (154–177)	.06
Initial respiratory rate per minute, median (IQR)	2675	48 (40–60)	2436	48 (40–60)	239	52 (40–62)	<.001
Initial oxygen saturation by pulse oximetry							.66
<90%	290	11	261	11	29	12	
90%–93.9%	424	16	384	16	40	17	
≥94%	1937	73	1771	73	166	71	
Initial retractions							.001
None	547	20	497	20	50	21	
Mild	1181	44	1091	44	90	37	
Moderate/severe	819	30	723	29	96	40	
Not documented/unknown	162	6	157	6	5	2	
Presence of apnea	174	6	165	7	9	4	.07
Presence of wheezing	1693	64	1554	65	139	59	.08
Inhaled bronchodilator use	1690	63	1531	63	159	66	.26
Systemic corticosteroid use	387	14	362	15	25	10	.06
Systemic antibiotic use	602	22	556	23	46	19	.19
Inpatient							
ICU admission	436	16	362	15	74	31	<.001
Continuous positive airway pressure and/or intubated	165	6	134	5	31	13	<.001
Presence of apnea	116	4	105	4	11	5	.85
Inhaled bronchodilator on first admission day	990	37	881	36	109	45	.007
Systemic corticosteroid use	434	16	398	16	36	15	.58
Systemic antibiotic use	302	30	249	29	53	32	.46
Hospital LOS, d, median (IQR)	2709	2 (1–4)	2468	2 (1–4)	241	3 (2–5)	<.001

results reported as odds ratios with 95% confidence intervals. All *P* values were 2-tailed, with *P* < .05 considered statistically significant.

RESULTS

HTS Use Among Children Hospitalized for Bronchiolitis

Overall, the median age of the 2709 children hospitalized for bronchiolitis was 3.7 months, and 60% were male (Table 1). In total, 241 (9%) children received HTS

during their ED or inpatient stay. Of these 241 children, most (69%) received HTS during their inpatient stay only, 15% received it in the ED only, and 16% received HTS in both settings.

From the 2008–2009 season to the 2011–2012 season, administration of HTS increased from 2% of patients to 27% of patients. In the 2012–2013 season, it dropped to 15% of patients and continued to drop to 11% of patients in the

2013–2014 season (*P* < .001, Fig 1A). From 2011 to 2012, most HTS was contributed by 3 out of 9 (33%) sites. From 2012 to 2013, it was contributed by 6 out of 17 (35%) sites, with 1 site using HTS on 91% (10 out of 11) of its enrolled patients. For sites with data for both seasons (2008–2009 and 2009–2010) of MARC-30, 31% (5 out of 16) of sites increased their use of HTS between the 2 enrollment seasons. The decrease in HTS use from 2013 to 2014 was largely due to 1 site




FIGURE 1 Trends in HTS use for bronchiolitis compared with relevant literature. A, HTS use among children hospitalized for bronchiolitis from 2008 to 2014. B, Number of PubMed publications evaluating HTS use for bronchiolitis from 2002 to 2014.

decreasing use from 58% to 29%. The 5 other sites with high use from 2012 to 2013 did not enroll patients in the 2013–2014 season. Although site

participation varied by enrollment season, the type of sites selected for participation (eg, academic medical centers) remained relatively consistent. Among the 6 sites that

participated in all 4 enrollment seasons, there was still a pronounced increase in the proportion of patients receiving HTS (2008–2009, 2%; 2009–2010, 6%;

2011–2012, 35%) and then a decline in use (2012–2013, 27%).

Of the 241 children receiving HTS, 31% were admitted to the ICU, compared with 15% of those who did not receive HTS ($P < .001$). Intubation was also more common among those receiving HTS compared with those who did not (8% vs 3%; $P = .001$). Of the patients who received HTS, 184 (77%) also received bronchodilator therapy: 75 (31%) only in the ED, 25 (10%) only in the inpatient setting, and 84 (35%) in both settings. The median LOS of patients who received HTS was 3 days (IQR, 2–5 days) vs a median of 2 days (IQR, 1–4 days, $P < .0001$) in other children.

In the multivariable model, children hospitalized during the 3 seasons from 2011 to 2014 were more likely to have received HTS compared with those enrolled in the 2008–2009 season (Table 2). Other factors associated with an increased likelihood of HTS use were ICU at birth, breathing problem started <1 day before arrival, parental rating of breathing problem as “severe,” previous use of inhaled bronchodilator before arrival, and respiratory syncytial virus (RSV) infection. Also, children ≥ 12 months (only included in MARC-30) were less likely to receive HTS than those <2 months of age.

Literature Search

Between 1998 and 2014, 69 total articles were published evaluating HTS effectiveness for bronchiolitis; 25 (36%) positive, 38 (55%) neutral, and 6 (9%) negative. Before 2010, 1 to 3 articles were published on the topic annually (Fig 1B). However, between years 2010 and 2011, a total of 22 relevant articles were published, of which 7 were positive, 13 were neutral, and 2 were negative. Between years 2012 and 2014, 36 additional articles were published, of which 12 were positive, 20 were neutral, and 4 were negative. All of the negative studies in this period came in 2014.

DISCUSSION

In our study, 9% of children hospitalized for bronchiolitis were treated with HTS. Among children who received HTS, indicators of

severe disease (ie, ICU admission, longer hospital LOS, <1 day of symptoms, and a parental rating of breathing problem as “severe”) were more common compared with children who did not receive HTS. HTS use varied over season and enrolling sites, with an initial increase from 2008 to 2012 and then a decline during the 2012–2014 seasons. This variation in HTS use over time remained robust after

adjustment for potential confounders in multivariable analyses.

Previous studies have noted wide practice variation in various treatments for bronchiolitis^{8–8,22–27}; however, national guidelines appear to be decreasing unnecessary testing and the use of ineffective treatments.^{25–28} Beyond supportive care, no treatment has been clearly established for children with

TABLE 2 Predictors of Receiving HTS among 2709 Infants Hospitalized for Bronchiolitis from 2008 to 2014

Characteristics	Odds Ratio	95% Confidence Interval		P
Enrollment season				
2008–2009	1.00	Reference	—	—
2009–2010	3.22	1.03	10.05	.04
2011–2012	14.17	4.52	44.38	<.001
2012–2013	7.33	2.87	18.74	<.001
2013–2014	5.02	2.20	11.46	<.001
Age at enrollment (mo)				
<2.0	1.00	Reference	—	—
2.0–5.9	1.02	0.66	1.58	.94
6.0–11.9	0.80	0.50	1.27	.34
≥ 12	0.39	0.19	0.79	.009
Sex				
Male	1.00	Reference	—	—
Female	0.95	0.72	1.24	.70
Race and/or ethnicity				
Non-Hispanic white	1.00	Reference	—	—
Non-Hispanic black or African American	0.69	0.43	1.11	.13
Hispanic	1.26	0.57	2.78	.58
Other	0.98	0.43	2.23	.97
Insurance				
Private	1.00	Reference	—	—
Public	0.64	0.40	1.02	.06
None	0.52	0.22	1.24	.14
Either or both parents with history of asthma	1.17	0.91	1.49	.22
Premature birth (≤ 37 wk)	1.06	0.79	1.43	.68
ICU or special care at birth	1.39	1.09	1.78	.008
Breathing problem began <1 d before ED visit	1.71	1.14	2.56	.01
Child discomfort in 24 h before ED visit				
None/mild	1.00	Reference	—	—
Moderate	1.09	0.78	1.52	.62
Severe	1.51	1.06	2.15	.02
Used inhaled bronchodilator for breathing problem before ED visit	1.62	1.18	2.24	.003
RSV	1.48	1.03	2.12	.03
Rhinovirus	1.06	0.72	1.55	.78

bronchiolitis. This study, however, is the first to show a rapid adoption in HTS use for bronchiolitis in parallel with literature supporting its novel use but before official sanctioning in national guidelines. The initial increase may have been due to more research publications describing HTS use in bronchiolitis patients. As time progressed, further research showed equivocal and inconsistent results for the efficacy of HTS,^{8,15,29} and its overall usage subsequently declined in parallel with our findings. The quick uptake is also likely related to its negligible side effect profile, especially when coadministered with bronchodilators,^{30–32} and the lack of other effective treatments. In this context, it is not surprising that sicker patients were more likely to receive HTS because clinicians would be more willing to try novel treatments in patients not improving after more established treatments failed. This can lead to confounding by severity in observational studies of medication effectiveness.³⁵

More recently, a systematic review reported on all available trials of HTS up to May 2015 and concluded that nebulized HTS is associated with reduced LOS among admitted infants and reduced hospital admissions when used in the outpatient setting.³⁴ However, a reanalysis of 2 large meta-analyses questions these results and shows that the improvement in LOS disappears when a single outlier study is removed and the results are controlled for day of illness at presentation.³⁵ Although further research is needed to clarify the role of HTS in the management of bronchiolitis, it is also important to study the use and uptake of new treatments among clinicians.

This study had several potential limitations. First, the uptake and usage of HTS by clinicians could be affected by a variety of factors that we did not measure. However, our observations paralleled usage trends seen in the literature on the treatment of bronchiolitis. Our literature search strategy was limited to a single database (PubMed) and our categorization of articles relied on the assessment of 1 author. In addition, this was an

observational study, which complicates the evaluation of the treatment effectiveness of HTS.³⁵ Also, our sample included only children hospitalized for bronchiolitis in academic medical centers and not all eligible children were enrolled, which may limit the generalizability of the results. Finally, not all study sites enrolled participants in all years; some findings might be due to site variability and not necessarily usage variability. The variability in site participation coincided with the largest changes observed in HTS use; therefore, these changes may be exaggerated. Despite these limitations, this study suggests that for severe bronchiolitis, providers are quick to trial novel therapies like HTS given the absence of other effective treatments.

CONCLUSIONS

The use of HTS in children hospitalized with bronchiolitis initially increased during the 2008–2012 seasons but then declined. The average usage over the course of our analysis was 9%, but it varied significantly across years and sites. Factors associated with HTS use included younger age, RSV infection, bronchodilator use before presentation, and need for NICU at birth; sicker patients also were more likely to receive HTS. We believe our findings are at least partially explained by the trends in HTS literature with a predominance of positive articles early on followed by an increase in negative and neutral articles after 2012. Although further research will likely clarify the role of HTS in the management of bronchiolitis, it is also important to study the use and uptake of new treatments by clinicians. By using new treatments before they are proven effective, especially when no clear alternatives are available, patients could be placed at unjustified risk for adverse events.

REFERENCES

1. U.S. Department of Health and Human Services, Health Resources and Services Administration, Maternal and Child Health Bureau. *Child Health USA 2011*. Rockville, MD: U.S. Department of Health and Human Services; 2011

2. Shay DK, Holman RC, Newman RD, Liu LL, Stout JW, Anderson LJ. Bronchiolitis-associated hospitalizations among US children, 1980-1996. *JAMA*. 1999;282(15):1440–1446
3. Hasegawa K, Tsugawa Y, Brown DF, Mansbach JM, Camargo CA Jr. Trends in bronchiolitis hospitalizations in the United States, 2000-2009. *Pediatrics*. 2013;132(1):28–36
4. Pelletier AJ, Mansbach JM, Camargo CA Jr. Direct medical costs of bronchiolitis hospitalizations in the United States. *Pediatrics*. 2006;118(6):2418–2423
5. Panitch HB. Respiratory syncytial virus bronchiolitis: supportive care and therapies designed to overcome airway obstruction. *Pediatr Infect Dis J*. 2003;22(suppl2):S83–S87; discussion S87–S88
6. Behrendt CE, Decker MD, Burch DJ, Watson PH; International RSV Study Group. International variation in the management of infants hospitalized with respiratory syncytial virus. *Eur J Pediatr*. 1998;157(3):215–220
7. Pint A, Taljaard M, McGahern C, et al. Management of bronchiolitis in community hospitals in Ontario: a multicentre cohort study. *Can J Emerg Med*. 2016 18(6):443–452
8. Willson DF, Horn SD, Hendley JO, Smout R, Gassaway J. Effect of practice variation on resource utilization in infants hospitalized for viral lower respiratory illness. *Pediatrics*. 2001; 108(4):851–855
9. Ralston SL, Lieberthal AS, Meissner HC, et al; American Academy of Pediatrics. Clinical practice guideline: the diagnosis, management, and prevention of bronchiolitis. *Pediatrics*. 2014;134(5). Available at: www.pediatrics.org/cgi/content/full/134/5/e1474
10. Zhang L, Mendoza-Sassi RA, Wainwright C, Klassen TP. Nebulised hypertonic saline solution for acute bronchiolitis in infants. *Cochrane Database Syst Rev*. 2013;(7):CD006458
11. Mandelberg A, Amirav I. Hypertonic saline or high volume normal saline for viral bronchiolitis: mechanisms and

- rationale. *Pediatr Pulmonol.* 2010;45(1):36–40
12. Kuzik BA, Al-Qadhi SA, Kent S, et al. Nebulized hypertonic saline in the treatment of viral bronchiolitis in infants. *J Pediatr.* 2007;151(3):266–270.e1
 13. Mandelberg A, Tal G, Witzling M, et al. Nebulized 3% hypertonic saline solution treatment in hospitalized infants with viral bronchiolitis. *Chest.* 2003;123(2):481–487
 14. Sarrell EM, Tal G, Witzling M, et al. Nebulized 3% hypertonic saline solution treatment in ambulatory children with viral bronchiolitis decreases symptoms. *Chest.* 2002;122(6):2015–2020
 15. Silver AH, Esteban-Cruciani N, Azzarone G, et al. 3% hypertonic saline versus normal saline in inpatient bronchiolitis: a randomized controlled trial. *Pediatrics.* 2015;136(6):1036–1043
 16. Mansbach JM, Piedra PA, Stevenson MD, et al; MARC-30 Investigators. Prospective multicenter study of children with bronchiolitis requiring mechanical ventilation. *Pediatrics.* 2012;130(3). Available at: www.pediatrics.org/cgi/content/full/130/3/e492
 17. Mansbach JM, Hasegawa K, Henke DM, et al. Respiratory syncytial virus and rhinovirus severe bronchiolitis are associated with distinct nasopharyngeal microbiota. *J Allergy Clin Immunol.* 2016;137(6):1909–1913.e4
 18. Beckham JD, Cadena A, Lin J, et al. Respiratory viral infections in patients with chronic, obstructive pulmonary disease. *J Infect.* 2005;50(4):322–330
 19. Knorr L, Fox JD, Tilley PA, Ahmed-Bentley J. Evaluation of real-time PCR for diagnosis of *Bordetella pertussis* infection. *BMC Infect Dis.* 2006;6:62
 20. Winchell JM, Thurman KA, Mitchell SL, Thacker WL, Fields BS. Evaluation of three real-time PCR assays for detection of *Mycoplasma pneumoniae* in an outbreak investigation. *J Clin Microbiol.* 2008;46(9):3116–3118
 21. Mansbach JM, Piedra PA, Teach SJ, et al; MARC-30 Investigators. Prospective multicenter study of viral etiology and hospital length of stay in children with severe bronchiolitis. *Arch Pediatr Adolesc Med.* 2012;166(8):700–706
 22. Christakis DA, Cowan CA, Garrison MM, Molteni R, Marcuse E, Zerr DM. Variation in inpatient diagnostic testing and management of bronchiolitis. *Pediatrics.* 2005;115(4):878–884
 23. Mansbach JM, Emond JA, Camargo CA Jr. Bronchiolitis in US emergency departments 1992 to 2000: epidemiology and practice variation. *Pediatr Emerg Care.* 2005;21(4):242–247
 24. Barben J, Kuehni CE, Trachsel D, Hammer J; Swiss Paediatric Respiratory Research Group. Management of acute bronchiolitis: can evidence based guidelines alter clinical practice? *Thorax.* 2008;63(12):1103–1109
 25. Johnson LW, Robles J, Hudgins A, Osburn S, Martin D, Thompson A. Management of bronchiolitis in the emergency department: impact of evidence-based guidelines? *Pediatrics.* 2013;131(suppl 1):S103–S109
 26. McCulloh RJ, Smitherman SE, Koehn KL, Alverson BK. Assessing the impact of national guidelines on the management of children hospitalized for acute bronchiolitis. *Pediatr Pulmonol.* 2014;49(7):688–694
 27. Parikh K, Hall M, Teach SJ. Bronchiolitis management before and after the AAP guidelines. *Pediatrics.* 2014;133(1). Available at: www.pediatrics.org/cgi/content/full/133/1/e1
 28. Sakellaropoulou A, Emporiadou M, Aivazis V, Mauromixalis J, Hatzistilianou M. Acute bronchiolitis in a paediatric emergency department of Northern Greece. Comparisons between two decades. *Arch Med Sci.* 2012;8(3):509–514
 29. Grewal S, Klassen TP. The tale of 2 trials: disentangling contradictory evidence on hypertonic saline for acute bronchiolitis. *JAMA Pediatr.* 2014;168(7):607–609
 30. Al-Ansari K, Sakran M, Davidson BL, El Sayyed R, Mahjoub H, Ibrahim K. Nebulized 5% or 3% hypertonic or 0.9% saline for treating acute bronchiolitis in infants. *J Pediatr.* 2010;157(4):630–634.e1
 31. Kuzik BA, Flavin MP, Kent S, et al. Effect of inhaled hypertonic saline on hospital admission rate in children with viral bronchiolitis: a randomized trial. *CJEM.* 2010;12(6):477–484
 32. Luo Z, Liu E, Luo J, et al. Nebulized hypertonic saline/salbutamol solution treatment in hospitalized children with mild to moderate bronchiolitis. *Pediatr Int.* 2010;52(2):199–202
 33. Clark S, Costantino T, Rudnitsky G, Camargo CA Jr. Observational study of intravenous versus oral corticosteroids for acute asthma: an example of confounding by severity. *Acad Emerg Med.* 2005;12(5):439–445
 34. Zhang L, Mendoza-Sassi RA, Klassen TP, Wainwright C. Nebulized hypertonic saline for acute bronchiolitis: a systematic review. *Pediatrics.* 2015;136(4):687–701
 35. Brooks CG, Harrison WN, Ralston SL. Association between hypertonic saline and hospital length of stay in acute viral bronchiolitis: a reanalysis of 2 meta-analyses. *JAMA Pediatr.* 2016;170(6):577–584

APPENDIX TABLE 1 Principal Investigators at the 16 Participating Sites in MARC-30

Eugene Mowad, MD	Akron Children's Hospital, Akron, OH
Jonathan Mansbach, MD, MPH	Boston Children's Hospital, Boston, MA
John Cheng, MD and Carlos Delgado, MD	Children's Healthcare of Atlanta at Egleston, Atlanta, GA
Lisa Zaoutis, MD	Children's Hospital of Philadelphia, Philadelphia, PA
M. Jason Sanders, MD	Children's Memorial Hermann Hospital, Houston, TX
Brian Pate, MD	Children's Mercy Hospital & Clinics, Kansas City, MO
Stephen Teach, MD, MPH	Children's National Medical Center, Washington, DC
Besh Barcega, MD	Loma Linda University Children's Hospital, Loma Linda, CA
Frank LoVecchio, DO	Maricopa Medical Center, Phoenix, AZ
Paul Hain, MD and Mark Riederer, MD	Monroe Carell Jr. Children's Hospital at Vanderbilt, Nashville, TN
Dorothy Damore, MD and Nikhil Shah, MD	New York Presbyterian Hospital, New York, NY
Michelle Stevenson, MD, MS	Norton Children's Hospital, Louisville, KY
Erin Stucky Fisher, MD	Rady Children's Hospital, San Diego, CA
Haitham Haddad, MD	Rainbow Babies & Children's Hospital, Cleveland, OH
Alan Schroeder, MD	Santa Clara Valley Medical Center, San Jose, CA
Charles Macias, MD, MPH	Texas Children's Hospital, Houston, TX

APPENDIX TABLE 2 Principal Investigators at the 17 Participating Sites in MARC-35

Amy Thompson, MD	Alfred I. duPont Hospital for Children, Wilmington, DE
Federico Laham, MD, MS	Arnold Palmer Hospital for Children, Orlando, FL
Jonathan Mansbach, MD, MPH	Boston Children's Hospital, Boston, MA
Vincent Wang, MD, MHA	Children's Hospital of Los Angeles, Los Angeles, CA
Michelle Dunn, MD	Children's Hospital of Philadelphia, Philadelphia, PA
Juan Celedon, MD, DrPH	Children's Hospital of Pittsburgh, Pittsburgh, PA
Michael Gomez, MD, MS	Children's Hospital at St. Francis, Tulsa, OK
Brian Pate, MD and Henry Puls, MD	Children's Mercy Hospital & Clinics, Kansas City, MO
Stephen Teach, MD, MPH	Children's National Medical Center, Washington, DC
Richard Strait, MD	Cincinnati Children's Hospital and Medical Center, Cincinnati, OH
Ilana Wanik, MD	Connecticut Children's Medical Center, Hartford, CT
Sujit Iyer, MD	Dell Children's Medical Center of Central Texas, Austin, TX
Ari R. Cohen, MD and Wayne Shreffler, MD, PhD	Massachusetts General Hospital, Boston, MA
Michelle Stevenson, MD, MS	Norton Children's Hospital, Louisville, KY
Anne Beasley, MD	Phoenix Children's Hospital, Phoenix, AZ
Thida Ong, MD	Seattle Children's Hospital, Seattle, WA
Charles Macias, MD, MPH	Texas Children's Hospital, Houston, TX

Multicenter Observational Study of the Use of Nebulized Hypertonic Saline to Treat Children Hospitalized for Bronchiolitis From 2008 to 2014

Joshua Davis, Amy D. Thompson, Jonathan M. Mansbach, Pedro A. Piedra, Kohei Hasegawa, Ashley F. Sullivan, Janice A. Espinola and Carlos A. Camargo, Jr

Hospital Pediatrics 2017;7;483

DOI: 10.1542/hpeds.2017-0002 originally published online July 31, 2017;

Updated Information & Services	including high resolution figures, can be found at: http://hosppeds.aappublications.org/content/7/8/483
References	This article cites 30 articles, 9 of which you can access for free at: http://hosppeds.aappublications.org/content/7/8/483.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Bronchiolitis http://classic.hosppeds.aappublications.org/cgi/collection/bronchiolitis_sub Pulmonology http://classic.hosppeds.aappublications.org/cgi/collection/pulmonology_sub RSV http://classic.hosppeds.aappublications.org/cgi/collection/rsv_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: https://shop.aap.org/licensing-permissions/
Reprints	Information about ordering reprints can be found online: http://classic.hosppeds.aappublications.org/content/reprints

Multicenter Observational Study of the Use of Nebulized Hypertonic Saline to Treat Children Hospitalized for Bronchiolitis From 2008 to 2014

Joshua Davis, Amy D. Thompson, Jonathan M. Mansbach, Pedro A. Piedra, Kohei Hasegawa, Ashley F. Sullivan, Janice A. Espinola and Carlos A. Camargo, Jr

Hospital Pediatrics 2017;7:483

DOI: 10.1542/hpeds.2017-0002 originally published online July 31, 2017;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://hosppeds.aappublications.org/content/7/8/483>

Hospital Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 2012. Hospital Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 2154-1663.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

