

Parent-Provider Miscommunications in Hospitalized Children

Alisa Khan, MD, MPH,^{ab} Stephanie L. Furtak, BA,^a Patrice Melvin, MPH,^c Jayne E. Rogers, RN, MSN,^d Mark A. Schuster, MD, PhD,^{ab} Christopher P. Landrián, MD, MPH^{ab,e}

ABSTRACT

BACKGROUND: Miscommunications lead to medical errors and suboptimal hospital experience. Parent-provider miscommunications are understudied.

OBJECTIVES: (1) Examine characteristics of parent-provider miscommunications about hospitalized children, (2) describe associations among parent-provider miscommunications, parent-reported errors, and hospital experience, and (3) compare parent and attending physician reports of parent-provider miscommunications.

METHODS: Prospective cohort study of 471 parents of 0- to 17-year-old medical inpatients in a pediatric hospital between May 1, 2013 and October 1, 2014. At discharge, parents reported parent-provider miscommunication and type (selecting all applicable responses), overall experience, and errors during hospitalization. During discharge billing, the attending physicians ($n = 52$) of a subset of patients ($n = 217$) also reported miscommunications, enabling comparison of parent and attending physician reports. We used logistic regression to examine characteristics of parent-reported miscommunications; McNemar's test to examine associations between miscommunications, errors, and top-box (eg, "excellent") experience; and generalized estimating equations to compare parent- and attending physician-reported miscommunication rates.

RESULTS: Parents completed 406 surveys (86.2% response rate). 15.3% of parents ($n = 62$) reported miscommunications. Parents of patients with nonpublic insurance (odds ratio: 1.99; 95% confidence interval: 1.03–3.85) and longer lengths of stay (odds ratio: 1.12; 95% confidence interval: 1.02–1.23) more commonly reported miscommunications. Parents reporting miscommunications were 5.3 times more likely to report errors and 78.6% less likely to report top-box overall experience ($P < .001$ for both). Among patients with both parent and attending physician surveys, 16.1% ($n = 35$) of parents and 3.7% ($n = 8$) of attending physicians reported miscommunications ($P < .001$). Both parents and attending physicians attributed miscommunications most often to family receipt of conflicting information.

CONCLUSIONS: Parent-provider miscommunications were associated with parent-reported errors and suboptimal hospital experience. Parents reported parent-provider miscommunications more often than attending physicians did.

FREE

^aDivision of General Pediatrics, ^bProgram for Patient Safety and Quality, and ^cDepartment of Nursing, Boston Children's Hospital, Boston, Massachusetts; ^dDepartment of Pediatrics, Harvard Medical School, Boston, Massachusetts; and ^eDivision of Sleep Medicine, Brigham and Women's Hospital, Boston, Massachusetts

www.hospitalpediatrics.org

DOI: <https://doi.org/10.1542/hpeds.2016-0190>

Copyright © 2017 by the American Academy of Pediatrics

Address correspondence to Alisa Khan, MD, MPH, Division of General Pediatrics, Department of Medicine, Boston Children's Hospital, 21 Autumn St, Room 200.2, Boston, MA 02215. E-mail: alisa.khan@childrens.harvard.edu

HOSPITAL PEDIATRICS (ISSN Numbers: Print, 2154-1663; Online, 2154-1671).

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Support for this work was provided by an Agency for Healthcare Research and Quality NRSA T32 HS000063 grant (trainee, Harvard-wide Pediatric Health Services Research Fellowship Program), an Agency for Healthcare Research and Quality K12HS022986 grant, an internal Boston Children's Hospital Program for Patient Safety and Quality grant, and a Boston Children's Hospital Taking on Tomorrow Innovation Award in Community/Patient Empowerment.

Miscommunications among health care team members are a leading cause of sentinel events¹—the most serious adverse events (AEs) occurring in hospitals. Although there is much research on communication between providers (physicians and nurses),^{2–6} less is known about the quality of communication between parents and providers during hospitalization.^{7–9}

Previous research suggests that parent-provider communication affects parents' hospital experience,^{10–15} which has implications for patient outcomes,^{16,17} hospital reimbursements,¹⁸ and hospital performance.¹⁹ Parent-provider communication also affects patient safety^{20–22} and malpractice risk.^{23,24}

We previously found that parent-provider communication breakdowns evidenced by lack of shared understanding about inpatient care plans occurred in 45.1% of parent-resident dyads.²⁵ We also found that quality of parent-provider nighttime communication was associated with overall hospital experience.¹⁰ We additionally found that parents provided unique information about hospital safety events, particularly communication-related ones.²²

However, little is known about types and predictors of parent-provider miscommunications. Therefore, in this study, we examined rates and types of parent-provider miscommunications and their associations with parent-reported errors and top-box experience in a cohort of hospitalized children. We also compared parent and attending physician reports of parent-provider miscommunications.

METHODS

Setting and Study Population

On a nightly basis, we surveyed parents of 4 randomly selected 0- to 17-year-old children within 1 day of anticipated discharge from 2 medical (nonsurgical, nonintensive care) inpatient units at a children's hospital from May 2013 to October 2014. We collected data as part of a multifaceted nighttime communication study.²⁶ We included patients from general pediatric, short-stay (straightforward patients with often, but not always, shorter

stays), and subspecialty (eg, hematology) services. We obtained parent verbal consent using a study information sheet. We administered parent written surveys on weekday (Monday to Thursday) evenings, collecting surveys that same evening or the following morning. We obtained patient sociodemographic and clinical characteristics via hospital administrative data. The hospital institutional review board approved the study.

Exclusions

Because of limited nighttime interpreter services, we excluded non-English speaking parents. We also excluded the parents of: patients spending <2 nights in the unit, patients boarding on the pediatric inpatient unit awaiting psychiatric placement, patients in state custody, and patients ≥ 18 years of age.

Parent Surveys

We collected data from a parent discharge experience survey¹⁰ developed with parent and survey methodologist input after piloting and cognitively testing the survey in our study units. Our survey included questions about (1) overall miscommunications, (2) overall parent-reported errors, and (3) overall (daytime and nighttime) hospital experience.

For miscommunications, parents responded to a closed (yes or no) item asking whether there were any communication problems between them and their child's care team (doctors and nurses) during hospitalization. Parents answering affirmatively further characterized communication problem(s) by selecting all applicable responses from 8 categories (conflicting information, delayed information, being upset by the manner of communication, not speaking directly with physician team, incorrect information, key information not provided, failing to understand the medical plan, and language barrier) and one write-in "other" option.

For errors, parents reported whether their child experienced any mistakes or any negative effects from mistakes during their overall hospitalization (2 items). Two physicians reviewed responses, classifying

them as errors, quality issues, or exclusions (Cohen's weighted $\kappa = 0.64$, agreement = 78%), and came to consensus to resolve discrepancies.²²

For hospital experience, parents rated the following constructs using a 5-point Likert scale: understanding of plan, quality of nighttime parent-provider communication, quality of nighttime physician-nurse communication, and overall (day and night) hospital experience (29 items, Cronbach $\alpha > 0.80$).

We previously published further details about the development and results of these parent-reported errors²² and overall experience¹⁰ items.

Attending Physician Surveys

We simultaneously surveyed attending physicians about miscommunications through an electronic quality improvement questionnaire incorporated into general pediatric and short-stay unit attending physician discharge billing (typically completed within 24 hours of discharge). Attending physicians reported miscommunications between each patient's parents and the care team (defined as nurses and physicians), using the same questions and categories as parents. Subspecialty and private (eg, outpatient pediatricians caring for their hospitalized patients) attending physicians did not answer these quality questions at discharge billing and, thus, did not provide miscommunication data.

Outcome Measures

Our primary outcome was parent-reported miscommunications. Secondary outcomes included parent-reported errors, top-box experience (described below), and attending physician-reported miscommunications.

Parent-Reported Errors and Top-Box Experience

We analyzed associations between parent responses to miscommunication items and their responses to error and experience items, respectively. After parent-reported errors were validated through 2-physician review,²² we dichotomized responses into those who reported an

error versus those who did not. We defined top-box experience as selecting the topmost score (eg, “excellent”) for all items comprising the survey’s overall experience construct. We dichotomized responses into those who did versus those who did not report top-box overall experience.

Sample Characteristics

We evaluated self-reported parent (age, sex, relationship to patient, race, ethnicity, education, income, and primary language spoken) and administrative-data–derived patient characteristics (age, sex, race, ethnicity, insurance, length of stay, and complex chronic condition [CCC] count). CCCs use *International Classification of Diseases, Ninth Revision, Clinical Modification* codes to identify medically complex children.²⁷

Statistical Analyses

We presented proportions of parent-reported miscommunications by using descriptive statistics. We compared patient and parent characteristics between parents who did and did not report miscommunications through χ^2 and Fisher’s exact tests for categorical variables. We used Wilcoxon-Mann-Whitney tests for medians and analysis of variance to compare means across groups. We analyzed parent- and patient-level predictors of miscommunications through multivariable logistic regression. A priori covariates of clinical significance with P values $\leq .20$ in bivariate analyses were considered for inclusion in the multivariable model. We used McNemar’s exact test to evaluate for associations between parent responses to the miscommunication items and their responses to the overall experience construct and error items. We compared parent and attending physician reporting of miscommunications by using a generalized estimating equation model to account for parent-provider clusters. A P value $< .05$ was considered statistically significant. We used REDCap (Vanderbilt University, Nashville, TN)²⁸ to manage study data and SAS 9.4 (SAS Institute, Cary, NC) for analyses.

RESULTS

Sample Characteristics

Overall, 98.9% ($n = 471$) of eligible parents consented for the study (Fig 1). Of these, 406 parents responded to the miscommunication item (86.2% response rate). Miscommunication survey data were available from attending physicians ($n = 52$) for a subset of these patients ($n = 217$), as described in the Methods section.

Patients were, on average, 6.1 (SD: 5.7) years old, predominantly non-Hispanic (69.5%) and nonpublicly insured (61.8%), with no CCCs (69.7%) and a median length of stay of 2.6 days (interquartile range [IQR]: 1.9–4.1) (Table 1). Parents were, on average, 36.8 years old (SD: 8.9) and predominantly female (67.7%), non-Hispanic (66.5%), primarily English-speaking (83.5%), and college-educated (65.5%), with 43.2% reporting an annual household income \geq \$50 000.

Parent-Reported Miscommunications

In the overall cohort of 406 parent respondents, 15.3% of parents ($n = 62$) reported miscommunications. Parents most commonly characterized miscommunications as receipt of conflicting (42%) or delayed (27%) information (Fig 2). See Table 2 for examples of parent-reported miscommunications from optional parent narrative comments.

In bivariate analysis, parents of patients with longer median lengths of stay (4.0 days [IQR 2.2–7.0] vs 2.3 days [IQR 1.9–3.9]; $P < .001$) and nonpublic insurance (74.2% vs 59.5%; $P = .03$) were more likely to report miscommunications. Miscommunications also significantly differed by parent ethnicity and income. Although Hispanic and low-income (\leq \$29 999/year) parents made up 15.8% and 21.4% of our overall study population respectively, Hispanic and low-income parents made up only 8.1% and 9.7% of those reporting miscommunications (Table 1).

In multivariable analysis, when controlling for patient ethnicity, CCC presence, and parent education, parents of nonpublicly insured patients (odds ratio [OR]: 1.99; 95% confidence interval [CI]: 1.03–3.85; $P = .04$)

and patients with longer lengths of stay (OR: 1.12; 95% CI: 1.02–1.23; $P = .02$) were more likely to report miscommunications (Table 3).

Parent-Reported Miscommunications Versus Errors and Top-Box Experience

We previously reported that 8.9% of parents reported safety incidents, of which 62.2% were determined to be errors,²² and that 42.5% of parents reported top-box overall experience.¹⁰ Parent-reported errors related to miscommunication included an operative procedure mistakenly discussed with a family after being filed in the wrong patient’s chart and an incorrect insulin pump rate corrected by a parent after day and night teams failed to communicate dose change during handoff. Parents reporting miscommunications were significantly more likely to report medical errors than parents not reporting miscommunications (OR: 5.32; 95% CI: 2.19–12.94; $P < .001$). Parents reporting miscommunications were also significantly less likely to report top-box overall experience than parents not reporting miscommunications (OR: 0.21; 95% CI: 0.11–0.44; $P < .001$).

Parent-Reported Versus Attending Physician-Reported Miscommunications

In the subset of 217 patients for whom paired parent and attending physician data were available, parents reported miscommunications significantly more often than attending physicians. Among this subset of patients, 16.1% of parents reported miscommunications, compared with only 3.7% of attending physicians ($P < .001$). After accounting for dyad clusters, parents were more likely to report miscommunications than attending physicians (OR: 5.02; 95% CI: 2.36–10.70; $P < .001$). Top categories of miscommunications among both parents and attending physicians included conflicting information provided to family (40% [$n = 14$] of parents; 37.5% [$n = 3$] of attending physicians), delayed provision of key information to family (34% [$n = 12$] of parents; 12% [$n = 1$] of attending physicians), and family being upset with communication (29% [$n = 10$] of parents;




FIGURE 1 Cohort diagram. General pediatric and short-stay attending physicians reported communication problems as part of a series of quality questions completed at the time of discharge billing.^a Subspecialty attending physicians were not asked these questions and thus did not contribute data for the paired parent-attending physician analysis.^b Patients with attending physician survey data (linked to billing codes) missing. For example, because of incorrect billing code or coverage by private attending physician.

25% [$n = 2$] of attending physicians) (Fig 2). Categories of miscommunications reported by parents but not identified by attending physicians included physician team not speaking directly with family (20% [$n = 7$]) and key information not provided to family (6% [$n = 2$]).

Among parent-attending physician dyads, 18.4% ($n = 40$) had a miscommunication reported by their parent and/or attending physician. Among these, 80% ($n = 32$) were reported only by the parent, 12% ($n = 5$) were reported only by the attending physician, and 8% ($n = 3$) were reported by both the parent and attending physician. Cases in which both the attending physician and the parent reported a miscommunication included provision of conflicting information to family and family being upset with communication.

DISCUSSION

We found that, in a cohort of children hospitalized at a pediatric hospital, more than 1 in 7 parents, but only 1 in 27 attending physicians, reported miscommunications between the family and the child's care team. Both parents and attending physicians most commonly

categorized miscommunications as due to conflicting information. Parents of children who were nonpublicly insured and had longer lengths of stay were more likely to report miscommunications. Parents reporting miscommunications were 5.3 times more likely to report errors and 78.6% less likely to report top-box overall experience scores than parents who did not report miscommunications.

Although multiple studies have shown that communication failures between providers are key contributors to patient safety events,^{2,29} less is known about the relationship between parent-provider miscommunications and patient safety. In our previous study of parent safety-reporting, parents were valid reporters, with nearly two-thirds of parent-reported events representing errors.²² Parents identified communication problems, including those occurring between parents and providers, as contributing factors in a number of these errors. Our finding in the current study that parents who reported miscommunications were 5.3 times more likely to report errors further substantiates the relationship between parent-provider

miscommunications and errors. Our results are also similar to those of a study revealing that adult patients with communication problems (eg, language barriers or disabilities) were 3 times more likely to experience preventable AEs than patients without communication problems.³⁰ In sum, our results suggest that poor communication between providers and patients and/or families may contribute to errors and AEs.

Beyond safety, suboptimal communication between providers and families can affect quality of care, including parent experience. Our group and others have previously found that parent-provider communication predicts parents' overall hospital experience.¹⁰⁻¹² Our previous study focused on the quality of parent-provider communication at night as a predictor of overall experience.¹⁰ The current study examines parent-provider miscommunications throughout hospitalization (day or night) as a predictor of overall experience, thereby reinforcing the link between parent-provider communication and overall experience. Like other studies, our study also identifies conflicting information as a common cause of parent-provider miscommunication.^{31,32} In addition to their effects on safety and hospital experience, miscommunications can lead to poor care transitions,^{11,33,34} postdischarge complications,³⁵⁻³⁷ readmissions,³⁸ and malpractice suits.^{24,39-41} In a recent study, for example, 7149 malpractice cases (leading to \$1.7 billion in losses) involved communication, with 55% of cases involving provider-patient communication failures.²³

Reasons for parent-provider hospital miscommunications may include parent/provider stress and sleep deprivation, which can impair comprehension and communication.⁴²⁻⁴⁵ Providers may lack time to adequately communicate with families,^{46,47} solicit questions,⁴⁸ or confirm understanding,⁴⁹ and fail to recognize the extent to which parents have questions or need clarification.^{50,51} Additionally, providers may use unfamiliar medical jargon⁵² and fail to actively engage

TABLE 1 Parent-Reported Miscommunications by Patient and Parent Characteristics

Characteristics	Overall, <i>n</i> = 406 (%)	≥1 Parent-Reported Miscommunication, <i>n</i> = 62 (%)	No Parent-Reported Miscommunication, <i>n</i> = 344 (%)	<i>P</i>
Patient				
Age, mean (SD), y	6.1 (5.7)	5.6 (5.9)	6.2 (5.7)	.42
Sex				
Female	203 (50.0)	32 (51.6)	171 (49.7)	.78
Male	203 (50.0)	30 (48.4)	173 (50.3)	
Race				
White	214 (52.7)	34 (54.8)	180 (52.3)	.91
African American	55 (13.5)	8 (12.9)	47 (13.7)	
Asian	6 (1.5)	1 (1.6)	5 (1.5)	
Other	82 (20.2)	10 (16.1)	72 (20.9)	
Multiracial	12 (3.0)	2 (3.2)	10 (2.9)	
Missing or unknown	37 (9.1)	7 (11.3)	30 (8.7)	
Ethnicity				
Hispanic	62 (15.3)	6 (9.7)	56 (16.3)	.20
Non-Hispanic	282 (69.5)	49 (79.0)	233 (67.7)	
Missing or unknown	62 (15.3)	7 (11.3)	55 (16.0)	
CCC^a presence				
No	301 (74.1)	42 (67.7)	259 (75.3)	.21
Yes	105 (25.9)	20 (32.3)	85 (24.7)	
Length of stay, median (IQR), d	2.6 (1.9–4.1)	4.0 (2.2–7.0)	2.3 (1.9–3.9)	<.001 ^c
Insurance				
Nonpublic	249 (61.8)	46 (74.2)	203 (59.5)	.03 ^e
Public	154 (38.2)	16 (25.8)	138 (40.5)	
Parent				
Age, mean (SD), y ^b	36.8 (8.9)	35.1 (9.2)	37.0 (8.9)	.17
Sex				
Female	275 (67.7)	38 (61.3)	237 (68.9)	.26
Male	62 (15.3)	9 (14.5)	53 (15.4)	
Missing	69 (17.0)	15 (24.2)	54 (15.7)	
Relationship to patient				
Parent	328 (80.8)	45 (72.6)	283 (82.3)	.15
Other	12 (3.0)	2 (3.2)	10 (2.9)	
Missing	66 (16.3)	15 (24.2)	51 (14.8)	
Race				
White	229 (56.4)	35 (56.5)	194 (56.4)	.37
African American	34 (8.4)	5 (8.1)	29 (8.4)	
Asian	14 (3.4)	2 (3.2)	12 (3.5)	
Other	34 (8.4)	3 (4.8)	31 (9.0)	
Multiracial	14 (3.4)	0 (0)	14 (4.1)	
Missing or unknown	81 (20.0)	17 (27.4)	64 (18.6)	
Ethnicity				
Hispanic	64 (15.8)	5 (8.1)	59 (17.2)	.04 ^e
Non-Hispanic	270 (66.5)	40 (64.5)	230 (66.9)	
Missing or unknown	72 (17.7)	17 (27.4)	55 (16.0)	
Parent primary language				
English	339 (83.5)	54 (87.1)	285 (82.8)	.41

TABLE 1 Continued

Characteristics	Overall, <i>n</i> = 406 (%)	≥1 Parent-Reported Miscommunication, <i>n</i> = 62 (%)	No Parent-Reported Miscommunication, <i>n</i> = 344 (%)	<i>P</i>
Other	67 (16.5)	8 (12.9)	59 (17.2)	
Education				
High school or less	68 (16.7)	6 (9.7)	62 (18.0)	.14
Any college	266 (65.5)	41 (66.1)	225 (65.4)	
Missing	72 (17.7)	15 (24.2)	57 (16.6)	
Annual household income				
≤\$29 999	87 (21.4)	6 (9.7)	81 (23.5)	.001 ^c
\$30 000–\$49 999	37 (9.1)	5 (8.1)	32 (9.3)	
\$50 000–\$74 999	27 (6.7)	12 (19.4)	15 (4.4)	
\$75 000–\$99 999	28 (6.9)	3 (4.8)	25 (7.3)	
≥\$100 000	120 (29.6)	18 (29.0)	102 (29.7)	
Missing	107 (26.4)	18 (29.0)	89 (25.9)	

^a The CCC system uses *International Classification of Diseases, Ninth Revision, Clinical Modification* codes to categorize medically complex children. It uses these codes to identify “any medical condition that can be reasonably expected to last at least 12 months (unless death intervenes) and to involve either several different organ systems or 1 organ system severely enough to require specialty pediatric care and probably some period of hospitalization in a tertiary care center.”²⁷

^b 18.7% of respondents did not report parent age.

^c Values are statistically significant.

families in care, including on family-centered rounds.^{53,54} Physicians may not explicitly share diagnostic or therapeutic uncertainty with families or explain that plans may be revised as clinical status evolves or consultants make recommendations. Parents may feel uncomfortable asking questions,^{48,55–57} particularly if there are perceived or actual gaps in language, power, or health literacy. Communication breakdowns between interprofessional team members (eg, physicians and nurses^{10,58} or primary and consulting services^{59,60}) may occur as the day progresses and be transmitted to parents.^{52,61,62}

Our finding that parents of patients with longer lengths of stay were more likely to report parent-provider miscommunications may reflect increased opportunities for communication breakdowns to occur with time (eg, due to more time for reporting and/or increased number of parent-provider interactions), increased illness complexity, or a combination of these factors. Additionally, providers may communicate less precisely or regularly with parents who are in the hospital longer if providers overestimate parents’ understanding of care plans and clinical trajectories.

Our finding that publicly insured patients have fewer miscommunications may reflect

underreporting by parents with fewer resources. Public insurance is a proxy for low socioeconomic status, which may also be associated with limited health literacy.⁶³ Limited health literacy is prevalent^{64,65} and associated with lower quality of physician-patient communication.⁶⁶ Although we did not directly measure health literacy in this study (education being an imperfect proxy⁶³), parents with limited health literacy may be less able to recognize miscommunications or feel less empowered to report them, leading to underreporting. We believe that, in reality, parents with lower health literacy are likely more susceptible to miscommunications because of limited comprehension of medical information and ability to ask appropriate clarifying questions. Further study of the relationship between health literacy and miscommunications is needed and is a topic we plan to explore in a future study.

Discrepancies between parent- and attending physician-reported miscommunication rates may relate to different definitions of, or thresholds for, reporting miscommunications among parents and attending physicians. For instance, attending physicians may overlook certain miscommunications that parents recognize. Additionally, attending physicians, who may be less involved in the day-to-day




FIGURE 2 Parent- and attending physician-reported miscommunications by type. Proportions and types of miscommunications reported by parents (*n* = 35) and by attending physicians (*n* = 8) among a subset of patients (*n* = 217) with paired parent and attending physician data are represented.

TABLE 2 Examples of Parent-Reported Miscommunications

Theme	Parent Narrative Comment
Delay in providing key information	"Sometimes things happen slower than they should."
Conflicting information	"Different info from different people." "Was told I no longer needed to [keep] wet diapers but later told I should keep for weighing (minor but inconsistent)."
The physician team did not speak directly to you	"[Doctor] never came to do exam." "I didn't know who my nighttime doctors were."
The way communication happened upset you	"[Felt] the doctors didn't value any input on my child's progress/care." "Nurses occasionally were not informed of changes in plans."
Other	"Treatment was discussed with attending during the day. Change in treatment by the attending not communicated (later by nurse)."

Representative quotations about miscommunications from parents taken from optional parent narrative responses.

more likely to respond and also hold higher expectations for optimal communication. Our ability to detect disparities by race and ethnicity and income was limited because there was a fair amount of missing data in these categories, although it does appear that there were some interesting trends worthy of further study.

Additionally, we did not collect attending physician demographic data. We also did not have access to perceptions of parent-provider miscommunications by subspecialty attending physicians, residents, or nurses, team members who might have additional important insights about miscommunications and who should be included in future studies. Furthermore, only the discharging attending physician

care of patients than residents and nurses, may not be privy to all miscommunications that occur with families. However, attending physicians are ultimately responsible for patient care and any safety, quality, or malpractice-related repercussions that may result from miscommunications on their care team. Therefore, it behooves them to be aware of miscommunications occurring between parents and the care team.

Discrepancies may also result from providers failing to recognize limited health literacy in families,^{49,67–69} providers (or parents) overestimating parent English comprehension, and providers overestimating the quality of communication with families. In a previous study in the same units, we found that 73.1% of providers reported they shared an understanding with parents about the care plan; however, when objectively rated, only 54.9% of provider-parent dyads were rated as having a shared understanding.²⁵ Additionally, multiple caregivers who do not communicate fully with one another may be present during the hospitalization.

Interventions to decrease hospital parent-provider miscommunications may include engaging patients and families in care through family-centered rounds^{53,54} and multifaceted,²⁰ multimodal communication techniques (eg, verbal and written); improving provider training in communication skills,^{70,71} cultural competency,^{72–74} and health literacy^{75,76}; enhancing interpreter use⁷⁷; and ensuring closed-loop communication through

strategies such as "teachback."^{49,78} Empowering parents (particularly those from disadvantaged backgrounds because of language, socioeconomic status, or health literacy barriers) to speak up when they have a question or need clarification may also reduce miscommunications. Although interventions to empower patients to speak up may lead to increased reports of miscommunications and errors, they will hopefully reduce their actual occurrence.

Given that both parents and attending physicians recognized conflicting information as a primary source of miscommunication, this may be a particular area of focus. Better engaging families in discussions, acknowledging uncertainty, being transparent that plans may change, and enhancing team communication (eg, between nurses and physicians, between day and night physicians, and between primary and consulting services)¹⁰ may help reduce the frequency at which parents receive conflicting information.

Limitations of our study include that it was conducted at a single-center institution among predominantly English-speaking, well-educated mothers of patients admitted for ≥ 2 nights. This limits generalizability as experiences of patients admitted for less time, from different backgrounds, or in other contexts may vary. For instance, it is likely that non-English speaking parents may experience more miscommunications.⁷⁷ Additionally, reporting bias may have affected our results if, for example, parents with higher socioeconomic status were

TABLE 3 Patient and Parent Factors Associated With Parent-Reported Miscommunications

	OR	95% CI	P
Patient			
Ethnicity			
Hispanic	0.67	0.29–1.82	.41
Missing or unknown	0.61	0.25–1.45	
Non-Hispanic	1.00	—	
CCC ^a presence			
Yes	1.20	0.65–2.23	.57
No	1.00	—	
Insurance			
Nonpublic	1.99	1.03–3.85	.04 ^b
Public	1.00	—	
Length of stay			
Per 2 d increase	1.12	1.02–1.23	.02 ^b
Parent			
Education			
Any college	1.32	0.51–3.42	.44
Missing	1.88	0.65–5.39	
High school or less	1.00	—	

—, not applicable.

^a The CCC system uses *International Classification of Diseases, Ninth Revision, Clinical Modification* codes to categorize medically complex children. It uses these codes to identify "any medical condition that can be reasonably expected to last at least 12 months (unless death intervenes) and to involve either several different organ systems or 1 organ system severely enough to require specialty pediatric care and probably some period of hospitalization in a tertiary care center."²⁷

^b Values are statistically significant.

was surveyed, not any previous attending physicians. Other limitations include our method and timing of recruitment: parent surveys were completed on weekdays only, which prevented us from detecting different, possibly higher, weekend miscommunication rates. Also, we recruited parents who were physically present at the bedside, who may have lower miscommunication rates than parents not present.

Future research should address the impact of parent-provider and provider-provider communication interventions on parent-provider miscommunications, as well as the relationship between parent-provider miscommunications and directly measured rates of errors and AEs. The roles of health literacy, socioeconomic, and language

barriers in parent-provider miscommunications also merit further study.

CONCLUSIONS

Miscommunications between parents and providers were common and underrecognized by attending physicians, although both parents and attending physicians recognized family receipt of conflicting information as a primary source of miscommunication. Parent-reported miscommunications were associated with parent-reported errors and suboptimal hospital experience. Improving communication between parents and providers has the potential to improve patient safety, patient and family experience, and malpractice risk.

Further research is needed to evaluate the effectiveness of interventions to improve parent-provider communication.

Acknowledgments

We wish to thank Thomas W. Mangione, PhD, for his assistance with reviewing the survey instrument; parent partners Brenda Allair, MA, and Katie Litterer, BA, for providing valuable parent perspectives; Fabienne Bourgeois, MD, MPH, for her assistance with implementing the attending physician quality survey and acquiring study data; Alexandra N. Mercer, BA, for providing revision and editing assistance for the manuscript; and all the families, physicians, nurses, and research assistants who participated in this study.

POTENTIAL CONFLICT OF INTEREST: Dr Landrigan has consulted with and holds equity in the I-PASS Institute, which seeks to train institutions in best handoff practices and aid in their implementation. Dr Landrigan has also served as a paid consultant to Virgin Pulse to help develop a Sleep and Health Program. He is supported in part by the Children's Hospital Association for his work as an Executive Council member of the Pediatric Research in Inpatient Settings (PRIS) network. In addition, Dr Landrigan has received monetary awards, honoraria, and travel reimbursement from multiple academic and professional organizations for teaching and consulting on sleep deprivation, physician performance, handoffs, and safety, and has served as an expert witness in cases regarding patient safety and sleep deprivation; the other authors have indicated they have no potential conflicts of interest to disclose.

The views in this article are solely those of the authors and do not necessarily represent the views of the Agency for Healthcare Research and Quality or of Boston Children's Hospital.

Dr Khan conceptualized and designed the study, obtained funding, acquired data, performed statistical analyses, analyzed and interpreted data, and drafted the initial manuscript; Ms Furtak participated in study design, tabulated articles, helped perform the literature review, and provided administrative support; Ms Melvin performed statistical analyses, and analyzed and interpreted data; Ms Rogers provided intellectual advice and guidance for the study and obtained funding; Dr Schuster provided intellectual advice and methodological guidance for the study; Dr Landrigan supervised the study, obtained funding, conceptualized and designed the study, and analyzed and interpreted data; and all authors critically reviewed and revised the manuscript for important intellectual content and approved the final manuscript as submitted.

REFERENCES

1. The Joint Commission Online. Sentinel event statistics released for 2014. Available at: www.jointcommission.org/assets/1/23/jconline_April_29_15.pdf. Accessed January 3, 2017
2. Starmer AJ, Spector ND, Srivastava R, et al; I-PASS Study Group. Changes in medical errors after implementation of a handoff program. *N Engl J Med*. 2014; 371(19):1803–1812
3. Horwitz LI, Moin T, Krumholz HM, Wang L, Bradley EH. Consequences of inadequate sign-out for patient care. *Arch Intern Med*. 2008;168(16):1755–1760
4. Arora V, Johnson J, Lovinger D, Humphrey HJ, Meltzer DO. Communication failures in patient sign-out and suggestions for improvement: a critical incident analysis. *Qual Saf Health Care*. 2005;14(6):401–407
5. Venkatesh AK, Curley D, Chang Y, Liu SW. Communication of vital signs at emergency department handoff: opportunities for improvement. *Ann Emerg Med*. 2015;66(2):125–130
6. Riesenber LA, Leitzsch J, Massucci JL, et al. Residents' and attending physicians' handoffs: a systematic review of the literature. *Acad Med*. 2009; 84(12):1775–1787
7. Moore JB, Kordick MF. Sources of conflict between families and health care professionals. *J Pediatr Oncol Nurs*. 2006;23(2):82–91
8. Fisher MJ, Broome ME. Parent-provider communication during hospitalization. *J Pediatr Nurs*. 2011;26(1):58–69
9. Azoulay E, Chevret S, Leleu G, et al. Half the families of intensive care unit patients experience inadequate communication with physicians. *Crit Care Med*. 2000;28(8):3044–3049
10. Khan A, Rogers JE, Melvin P, et al. Physician and nurse nighttime communication and parents' hospital experience. *Pediatrics*. 2015;136(5). Available at: www.pediatrics.org/cgi/content/full/136/5/e1249
11. Co JPT, Ferris TG, Marino BL, Homer CJ, Perrin JM. Are hospital characteristics associated with parental views of

- pediatric inpatient care quality? *Pediatrics*. 2003;111(2):308–314
12. Homer CJ, Marino B, Cleary PD, et al. Quality of care at a children's hospital: the parent's perspective. *Arch Pediatr Adolesc Med*. 1999;153(11):1123–1129
 13. Nobile C, Drotar D. Research on the quality of parent-provider communication in pediatric care: implications and recommendations. *J Dev Behav Pediatr*. 2003;24(4):279–290
 14. Hart CN, Kelleher KJ, Drotar D, Scholle SH. Parent-provider communication and parental satisfaction with care of children with psychosocial problems. *Patient Educ Couns*. 2007;68(2):179–185
 15. Ammentorp J, Mainz J, Sabroe S. Determinants of priorities and satisfaction in pediatric care. *Pediatr Nurs*. 2006;32(4):333–340,348
 16. Kane RL, Maciejewski M, Finch M. The relationship of patient satisfaction with care and clinical outcomes. *Med Care*. 1997;35(7):714–730
 17. Covinsky KE, Rosenthal GE, Chren MM, et al. The relation between health status changes and patient satisfaction in older hospitalized medical patients. *J Gen Intern Med*. 1998;13(4):223–229
 18. Centers for Medicare & Medicaid Services. Fact sheets: CMS issues final rule for first year of hospital value-based purchasing program. Available at: <https://www.cms.gov/Newsroom/MediaReleaseDatabase/Fact-sheets/2011-Fact-sheets-items/2011-04-29.html>. Accessed October 22, 2016
 19. Centers for Medicare & Medicaid Services. HCAHPS: patients' perspectives of care survey. Available at: www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/HospitalQualityInits/HospitalHCAHPS.html. Accessed October 31, 2016
 20. Fassier T, Azoulay E. Conflicts and communication gaps in the intensive care unit. *Curr Opin Crit Care*. 2010;16(6):654–665
 21. Azoulay E, Timsit J-F, Sprung CL, et al; Conflicus Study Investigators and for the Ethics Section of the European Society of Intensive Care Medicine. Prevalence and factors of intensive care unit conflicts: the conflicus study. *Am J Respir Crit Care Med*. 2009;180(9):853–860
 22. Khan A, Furtak SL, Melvin P, Rogers JE, Schuster MA, Landrigan CP. Parent-reported errors and adverse events in hospitalized children. *JAMA Pediatr*. 2016;170(4):e154608
 23. CRICO Strategies. *Malpractice Risks in Communication Failures: 2015 Annual Benchmarking Report*. Boston, MA: The Risk Management Foundation of Harvard Medical Institutions Incorporated; 2015. Available at: <https://www.rmf.harvard.edu/cbsreport>
 24. Beckman HB, Markakis KM, Suchman AL, Frankel RM. The doctor-patient relationship and malpractice. Lessons from plaintiff depositions. *Arch Intern Med*. 1994;154(12):1365–1370
 25. Khan A, Rogers JE, Forster CS, Furtak SL, Schuster MA, Landrigan CP. Communication and shared understanding between parents and resident-physicians at night. *Hosp Pediatr*. 2016;6(6):319–329
 26. Khan A, Baird J, Rogers JE, et al. Parent and provider experience and shared understanding after a family-centered nighttime communication intervention. *Acad Pediatr*. 2017;17(4):389–402
 27. Feudtner C, Christakis DA, Connell FA. Pediatric deaths attributable to complex chronic conditions: a population-based study of Washington State, 1980-1997. *Pediatrics*. 2000;106(suppl 1, 1 pt 2):205–209
 28. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377–381
 29. Kohn L, Corrigan J, Donaldson MS; Institute of Medicine. To err is human: building a safer health system. Available at: <http://iom.nationalacademies.org/~media/Files/Report%20Files/1999/To-Err-is-Human/To%20Err%20is%20Human%201999%20report%20brief.pdf>. Accessed October 31, 2016
 30. Bartlett G, Blais R, Tamblyn R, Clermont RJ, MacGibbon B. Impact of patient communication problems on the risk of preventable adverse events in acute care settings. *CMAJ*. 2008;178(12):1555–1562
 31. Clarke JN, Fletcher P. Communication issues faced by parents who have a child diagnosed with cancer. *J Pediatr Oncol Nurs*. 2003;20(4):175–191
 32. Mercer M, Ritchie JA. Home community cancer care: parents' perspectives. *J Pediatr Nurs*. 1997;12(3):133–141
 33. Kripalani S, Jackson AT, Schnipper JL, Coleman EA. Promoting effective transitions of care at hospital discharge: a review of key issues for hospitalists. *J Hosp Med*. 2007;2(5):314–323
 34. Sellick S, Desa V, MacDonald K. The impact of shared care on children with cancer and their families. *Paediatr Child Health*. 1996;1(3):211–215
 35. Tsilimingras D, Bates DW. Addressing postdischarge adverse events: a neglected area. *Jt Comm J Qual Patient Saf*. 2008;34(2):85–97
 36. Bernstein HH, Spino C, Lalama CM, Finch SA, Wasserman RC, McCormick MC. Unreadiness for postpartum discharge following healthy term pregnancy: impact on health care use and outcomes. *Acad Pediatr*. 2013;13(1):27–39
 37. Knight DA, Thompson D, Mathie E, Dickinson A. 'Seamless care? Just a list would have helped!' Older people and their carer's experiences of support with medication on discharge home from hospital. *Health Expect*. 2013;16(3):277–291
 38. Brittan M, Albright K, Cifuentes M, Jimenez-Zambrano A, Kempe A. Parent and provider perspectives on pediatric readmissions: what can we learn about readiness for discharge? *Hosp Pediatr*. 2015;5(11):559–565
 39. Patel AJK, Morrison CM. Opportunities to reduce plastic surgery claims through

- an analysis of complaints data. *J Plast Reconstr Aesthet Surg*. 2013;66(4): 455–459
40. Schaffer AC, Puopolo AL, Raman S, Kachalia A. Liability impact of the hospitalist model of care. *J Hosp Med*. 2014;9(12):750–755
 41. Whang JS, Baker SR, Patel R, Luk L, Castro A III. The causes of medical malpractice suits against radiologists in the United States. *Radiology*. 2013;266(2): 548–554
 42. Papp KK, Stoller EP, Sage P, et al. The effects of sleep loss and fatigue on resident-physicians: a multi-institutional, mixed-method study. *Acad Med*. 2004; 79(5):394–406
 43. Fostick L, Babkoff H, Zukerman G. Effect of 24 hours of sleep deprivation on auditory and linguistic perception: a comparison among young controls, sleep-deprived participants, dyslexic readers, and aging adults. *J Speech Lang Hear Res*. 2014;57(3):1078–1088
 44. van der Helm E, Gujar N, Walker MP. Sleep deprivation impairs the accurate recognition of human emotions. *Sleep*. 2010;33(3):335–342
 45. McLoone JK, Wakefield CE, Yoong SL, Cohn RJ. Parental sleep experiences on the pediatric oncology ward. *Support Care Cancer*. 2013;21(2):557–564
 46. McPherson G, Jefferson R, Kissoon N, Kwong L, Rasmussen K. Toward the inclusion of parents on pediatric critical care unit rounds. *Pediatr Crit Care Med*. 2011;12(6):e255–e261
 47. Turner T, Cull WL, Bayldon B, et al. Pediatricians and health literacy: descriptive results from a national survey. *Pediatrics*. 2009;124(suppl 3): S299–S305
 48. Katz MG, Jacobson TA, Veledar E, Kripalani S. Patient literacy and question-asking behavior during the medical encounter: a mixed-methods analysis. *J Gen Intern Med*. 2007;22(6): 782–786
 49. Schillinger D, Piette J, Grumbach K, et al. Closing the loop: physician communication with diabetic patients who have low health literacy. *Arch Intern Med*. 2003;163(1):83–90
 50. Howard T, Jacobson KL, Kripalani S. Doctor talk: physicians' use of clear verbal communication. *J Health Commun*. 2013;18(8):991–1001
 51. Collins TC, Clark JA, Petersen LA, Kressin NR. Racial differences in how patients perceive physician communication regarding cardiac testing. *Med Care*. 2002;40(suppl 1):I27–I34
 52. White C, Del Rey JG. Decreasing adverse events through night talks: an interdisciplinary, hospital-based quality improvement project. *Perm J*. 2009; 13(4):16–22
 53. Mittal V. Family-centered rounds. *Pediatr Clin North Am*. 2014;61(4):663–670
 54. Subramony A, Hametz PA, Balmer D. Family-centered rounds in theory and practice: an ethnographic case study. *Acad Pediatr*. 2014;14(2):200–206
 55. Arnold CL, Coran JJ, Hagen MG. Revisiting patient communication training: an updated needs assessment and the AGENDA model. *Patient Educ Couns*. 2012;88(3):399–405
 56. Parikh NS, Parker RM, Nurss JR, Baker DW, Williams MV. Shame and health literacy: the unspoken connection. *Patient Educ Couns*. 1996;27(1):33–39
 57. Street RL. Communicative styles and adaptations in physician-parent consultations. *Soc Sci Med*. 1992;34(10): 1155–1163
 58. Akl EA, Bais A, Rich E, Izzo J, Grant BJB, Schünemann HJ. Brief report: internal medicine residents', attendings', and nurses' perceptions of the night float system. *J Gen Intern Med*. 2006;21(5): 494–497
 59. Schmocker RK, Holden SE, Vang X, et al. The number of inpatient consultations is negatively correlated with patient satisfaction in patients with prolonged hospital stays. *Am J Surg*. 2016;212(2): 282–288
 60. Stevens JP, Johansson AC, Schonberg MA, Howell MD. Elements of a high-quality inpatient consultation in the intensive care unit. A qualitative study. *Ann Am Thorac Soc*. 2013;10(3):220–227
 61. Gonzalo JD, Moser E, Lehman E, Kuperman E. Quality and safety during the off hours in medicine units: a mixed methods study of front-line provider perspectives. *J Hosp Med*. 2014;9(12): 756–763
 62. Trontell MC, Carson JL, Taragin MI, Duff A. The impact of the night float system on internal medicine residency programs. *J Gen Intern Med*. 1991;6(5): 445–449
 63. Martin LT, Ruder T, Escarce JJ, et al. Developing predictive models of health literacy. *J Gen Intern Med*. 2009;24(11): 1211–1216
 64. Kutner M, Greenburg E, Jin Y, Paulsen C; National Center for Education Statistics. The health literacy of America's adults: results from the 2003 National Assessment of Adult Literacy. Available at: <http://nces.ed.gov/pubs2006/2006483.pdf>. Accessed October 24, 2016
 65. Weiss BD. Removing barriers to better, safer care: health literacy and patient safety: helping patients understand. AMA Foundation. Available at: www.ama-assn.org/ezp-prod1.hul.harvard.edu/resources/doc/ama-foundation/healthlitclinicians.pdf. Accessed July 18, 2017
 66. Kripalani S, Jacobson TA, Mugalla IC, Cawthon CR, Niesner KJ, Vaccarino V. Health literacy and the quality of physician-patient communication during hospitalization. *J Hosp Med*. 2010;5(5): 269–275
 67. Kelly PA, Haidet P. Physician overestimation of patient literacy: a potential source of health care disparities. *Patient Educ Couns*. 2007; 66(1):119–122
 68. Safer RS, Keenan J. Health literacy: the gap between physicians and patients. *Am Fam Physician*. 2005;72(3):463–468
 69. Vargas CR, Chuang DJ, Lee BT. Assessment of patient health literacy: a national survey of plastic surgeons. *Plast Reconstr Surg*. 2014;134(6): 1405–1414

70. Cegala DJ, Lenzmeier Broz S. Physician communication skills training: a review of theoretical backgrounds, objectives and skills. *Med Educ.* 2002;36(11):1004–1016
71. Weiss S, Goldlust E, Vaucher YE. Improving parent satisfaction: an intervention to increase neonatal parent-provider communication. *J Perinatol.* 2010;30(6):425–430
72. Lapinski J, Colonna C, Sexton P, Richard M. American sign language and deaf culture competency of osteopathic medical students. *Am Ann Deaf.* 2015;160(1):36–47
73. Turner DA, Fleming GM, Winkler M, et al; Education in Pediatric Intensive Care Investigators. Professionalism and communication education in pediatric critical care medicine: the learner perspective. *Acad Pediatr.* 2015;15(4):380–385
74. Lee AL, Mader EM, Morley CP. Teaching cross-cultural communication skills online: a multi-method evaluation. *Fam Med.* 2015;47(4):302–308
75. Coleman CA, Fromer A. A health literacy training intervention for physicians and other health professionals. *Fam Med.* 2015;47(5):388–392
76. Mackert M, Ball J, Lopez N. Health literacy awareness training for healthcare workers: improving knowledge and intentions to use clear communication techniques. *Patient Educ Couns.* 2011;85(3):e225–e228
77. Flores G, Abreu M, Barone CP, Bachur R, Lin H. Errors of medical interpretation and their potential clinical consequences: a comparison of professional versus ad hoc versus no interpreters. *Ann Emerg Med.* 2012;60(5):545–553
78. Griffey RT, Shin N, Jones S, et al. The impact of teach-back on comprehension of discharge instructions and satisfaction among emergency patients with limited health literacy: A randomized, controlled study. *J Commun Healthc.* 2015;8(1):10–21

Parent-Provider Miscommunications in Hospitalized Children

Alisa Khan, Stephannie L. Furtak, Patrice Melvin, Jayne E. Rogers, Mark A. Schuster
and Christopher P. Landrigan

Hospital Pediatrics 2017;7;505

DOI: 10.1542/hpeds.2016-0190 originally published online August 2, 2017;

Updated Information & Services	including high resolution figures, can be found at: http://hosppeds.aappublications.org/content/7/9/505
References	This article cites 70 articles, 7 of which you can access for free at: http://hosppeds.aappublications.org/content/7/9/505#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Administration/Practice Management http://hosppeds.aappublications.org/cgi/collection/administration:practice_management_sub Hospital Medicine http://hosppeds.aappublications.org/cgi/collection/hospital_medicine_sub Interpersonal & Communication Skills http://hosppeds.aappublications.org/cgi/collection/interpersonal_-_communication_skills_sub Patient Education/Patient Safety/Public Education http://hosppeds.aappublications.org/cgi/collection/patient_education:patient_safety:public_education_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://hosppeds.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://hosppeds.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™





Parent-Provider Miscommunications in Hospitalized Children

Alisa Khan, Stephannie L. Furtak, Patrice Melvin, Jayne E. Rogers, Mark A. Schuster
and Christopher P. Landrigan

Hospital Pediatrics 2017;7;505

DOI: 10.1542/hpeds.2016-0190 originally published online August 2, 2017;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://hosppeds.aappublications.org/content/7/9/505>

Hospital Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 2012. Hospital Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 2154-1663.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

