

To Spinal Tap or Not To Spinal Tap, That Is the Question

Thomas J. Lee, MD,^a Paul L. Aronson, MD^{a,b}

CASE PRESENTATION

Marcia is a 35-day-old, previously healthy girl who presents to the emergency department (ED) for “feeling warm.” Marcia has been feeding well with a normal amount of wet diapers. There are no other symptoms, including no fussiness, rash, rhinorrhea, diarrhea, vomiting, cough, or respiratory distress. Marcia had been alert and happy at home, but this evening she felt warm to her mother's touch. Marcia's birth history was unremarkable; she was born at term via vaginal delivery without complication and after appropriate prenatal care. She enjoys a diet of breast milk and has been growing and developing well.

In the ED, Marcia's vital signs are as follows: rectal temperature is 101.9°F, heart rate is 140 beats per minute, respiratory rate is 45 breaths per minute, blood pressure is 76/51, and pulse oximetry is 99% on room air. She is well appearing with no significant findings on physical examination. A complete blood count, procalcitonin, blood culture, urinalysis, and catheterized urine culture are obtained and sent for analysis. Her laboratory values are notable for a peripheral white blood cell count of 10 000 cells/ μ L with an absolute neutrophil count of 7000 cells/ μ L and no band cells. Her procalcitonin is 0.25 ng/mL, and the urinalysis is negative for leukocyte esterase and nitrites with no leukocytes on microscopy. A lumbar puncture is performed, and on the third attempt, cerebrospinal fluid (CSF) is obtained with results as follows: white blood cell count is 20 cells/ μ L, red blood cell count is 14 000 cells/ μ L, protein is 71 mg/dL, and glucose is 60 mg/dL. Because of the CSF pleocytosis, Marcia is admitted to the general pediatrics floor after receipt of ampicillin and ceftriaxone at meningitic dosing. She is hospitalized on intravenous antibiotics for 36 hours. During the hospital stay, her intravenous line infiltrates after her second dose of antibiotics, which leads to the placement of a second intravenous line after multiple attempts. Marcia is ultimately discharged from the hospital with outpatient follow-up and instructions to return if she becomes ill appearing. Her urine, blood, and CSF culture results remain negative for 5 days.

WHAT IS THE VALUE OF CSF TESTING IN FEBRILE INFANTS?

Evaluation and management of the febrile infant <90 days of age is a common scenario for general pediatricians, emergency medicine physicians, and hospitalists.¹ Although the prevalence of serious bacterial infection (SBI) in febrile infants is ~10%,^{2,3} most of these infections are urinary tract infections. Bacterial meningitis is rare, with an overall prevalence of ~0.6% in febrile infants,⁴ and is even lower among well-appearing febrile infants >28 days of

www.hospitalpediatrics.org

DOI:https://doi.org/10.1542/hpeds.2017-0207

Copyright © 2018 by the American Academy of Pediatrics

Address correspondence to Paul L. Aronson, MD, Department of Pediatrics, Section of Pediatric Emergency Medicine, Yale School of Medicine, 100 York St, Suite 1F, New Haven, CT 06511. E-mail: paul.aronson@yale.edu

HOSPITAL PEDIATRICS (ISSN Numbers: Print, 2154-1663; Online, 2154-1671).

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Supported by the Clinical and Translational Science Awards Program (grant KL2 TR001862) (Aronson) from the National Center for Advancing Translational Science, a component of the National Institutes of Health (NIH). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

Dr Lee conceptualized and designed the study and drafted the initial manuscript; Dr Aronson conceptualized and designed the study and critically revised the manuscript for important intellectual content; and all authors approved the final manuscript as submitted.



^aDepartments of
Pediatrics and
^bEmergency Medicine,
Yale School of Medicine,
New Haven, Connecticut

age.^{1,5,6} This low prevalence of meningitis coupled with the risks and costs of lumbar puncture and resultant hospitalization warrants the ongoing discussion about the value of CSF testing in a well-appearing febrile infant.

WHAT ARE CURRENT PRACTICE PATTERNS?

With numerous risk stratification criteria and algorithms available for reference,³ the percentage of febrile infants who undergo lumbar puncture varies widely across institutions and clinicians.^{1,7} However, the percentage of febrile infants who undergo lumbar puncture is lower for infants evaluated in the office setting compared with the ED.^{1,5,7} By using clinical judgment when deciding to perform invasive testing, primary care practitioners identify a similar proportion of infants with bacterial meningitis compared with those using a more strict adherence to established criteria. In a study of over 3000 febrile infants in an office setting, only 40% of infants underwent lumbar puncture. Among the 63 infants with bacteremia and/or bacterial meningitis in the cohort, there was only 1 infant with a delayed diagnosis of bacterial meningitis, and this infant was reported to have done well.⁵ In a recent study at Kaiser Permanente, researchers found that up to one-third of febrile infants did not have any bacterial cultures obtained, including 24% of infants 7 to 28 days old, without associated poor outcomes or increased return visits. Furthermore, there was no delayed identification of bacteremia or bacterial meningitis in the cohort.¹ These outpatient investigations are in contrast to studies of management in the pediatric ED that have revealed that neonates ≤ 28 days of age routinely undergo lumbar puncture and hospitalization, whereas $\sim 50\%$ of infants 29 to 56 days have CSF testing performed.⁷⁻⁹

ARE THERE RISKS WITH ROUTINE VERSUS SELECTIVE CSF TESTING?

One can argue that febrile infants managed in the primary care office are fundamentally different from those evaluated in the ED,

particularly in regard to reliable follow-up. Although this may partially explain the differences in management, the potential downsides to strategies that use routine versus more selective lumbar puncture must also be identified. For well-appearing febrile infants who are otherwise low risk for an SBI, the risk of not performing a lumbar puncture is a delay in diagnosis of bacterial meningitis. The following risk stratification algorithms do not include CSF testing: the Rochester criteria,¹⁰ the modified Philadelphia protocol,¹¹ and the Step-by-Step approach.⁶ Although these algorithms differ in their age cutoffs and in the specific laboratory tests used to define low risk, infants who are older than 28 days, well appearing, and have reassuring urine and blood test results have a risk of meningitis as low as 0%; however, the small number of cases of meningitis make this risk estimate imprecise.^{6,10,11} In addition, although meningitis is associated with mortality and neurologic morbidity in young infants,^{12,13} the few infants who were deemed low risk through stratification strategies but were ultimately diagnosed with meningitis have had reportedly good outcomes.³ However, there is a paucity of data on these infants given the very small number of infants in the published literature that were initially stratified as low risk and then subsequently diagnosed with meningitis; therefore, the absolute risk of an adverse outcome is unclear.³ What is clear is that the prevalence of bacterial meningitis is very low among these otherwise low-risk infants.

The reverse side of the coin is the potential downside to routine performance of lumbar punctures. First, the results of the CSF testing can be ambiguous. As in our patient, traumatic lumbar punctures can cloud the use of risk stratification criteria. Approximately 14% to 18% of lumbar punctures attempted are either traumatic or unsuccessful.^{14,15} Additionally, certain factors may increase the probability of a traumatic or unsuccessful lumbar puncture in a febrile infant, such as performance of the procedure by a less experienced provider, absence of local anesthetic use, and advancement of the spinal needle with the stylet in place.¹⁶ Given the inherent

difficulty in the interpretation of CSF results, traumatic lumbar punctures lead to more hospitalizations,¹⁴ which in turn result in higher costs.⁸ This increased resource use occurs despite a similar prevalence of bacterial meningitis in these infants compared with those with nontraumatic lumbar punctures.^{14,15} In addition to unnecessary hospitalization, traumatic lumbar punctures may also lead to repeat procedures, which amplifies the risk of iatrogenic morbidity. Furthermore, a higher proportion of positive CSF culture results represents contaminants rather than true pathogens,¹⁷ which can result in prolonged hospital stay and longer duration of intravenous antibiotics.

It is also important to note the potential “downstream” effects of lumbar puncture and hospitalization. The performance of lumbar puncture with subsequent hospitalization is highly stressful and anxiety provoking for parents.¹⁸ Additionally, hospitalized infants may be exposed to nosocomial infections and iatrogenic complications that carry risks and associated costs. Specifically, up to 20% of febrile infants have been reported to experience a complication during the hospitalization, which range from the infiltrated intravenous line in our case to uncommon but more serious adverse events.^{3,19} Lastly, hospitalized infants often receive empirical intravenous antibiotic therapy,⁷ and antibiotic use in early infancy is linked to asthma and pediatric obesity, each with potential morbidity.²⁰⁻²²

RISK TOLERANCE

Weighing all of these factors, the decision to perform a lumbar puncture and hospitalize a febrile infant is often based on tolerance of risk. Given the low prevalence of bacterial meningitis in low-risk febrile infants, particularly by using newer algorithms such as the Step-by-Step approach,⁶ many well-appearing febrile infants >28 days of age do not require a lumbar puncture and hospitalization. The decision on whether to perform a lumbar puncture should take into account not just the physician's tolerance for risk and confidence in the examination of young infants but also the risk tolerance of the

parents through a shared decision-making process.²³ The decision will also depend on the ability to ensure close follow-up and, therefore, engagement of the primary care physician may also be of benefit. Implementation of evidence-based protocols for management of febrile infants, such as the care process model implemented in the Intermountain Healthcare system, can also improve resource use and diagnosis of SBI while simultaneously lowering costs.²⁴ Further studies are still needed to assess outcomes for infants who experience a delayed diagnosis of SBI, particularly meningitis, as well as to evaluate the outpatient management without lumbar puncture for low-risk neonates 7 to 28 days of age.³ Ultimately, outpatient observation without lumbar puncture for low-risk febrile infants is a cost-effective, value-based management strategy for this patient population.

REFERENCES

- Greenhow TL, Hung YY, Pantell RH. Management and outcomes of previously healthy, full-term, febrile infants ages 7 to 90 days. *Pediatrics*. 2016;138(6):e20160270
- Huppler AR, Eickhoff JC, Wald ER. Performance of low-risk criteria in the evaluation of young infants with fever: review of the literature. *Pediatrics*. 2010;125(2):228–233
- Hui C, Neto G, Tsertsvadze A, et al. Diagnosis and management of febrile infants (0-3 months). *Evid Rep Technol Assess (Full Rep)*. 2012;205:1–297
- Cruz AT, Mahajan P, Bonsu BK, et al; Febrile Infant Working Group of the Pediatric Emergency Care Applied Research Network. Accuracy of complete blood cell counts to identify febrile infants 60 days or younger with invasive bacterial infections. *JAMA Pediatr*. 2017;171(11):e172927
- Pantell RH, Newman TB, Bernzweig J, et al. Management and outcomes of care of fever in early infancy. *JAMA*. 2004;291(10):1203–1212
- Gomez B, Mintegi S, Bressan S, Da Dalt L, Gervais A, Lacroix L; European Group for Validation of the Step-by-Step Approach. Validation of the “Step-by-Step” approach in the management of young febrile infants. *Pediatrics*. 2016;138(2):e20154381
- Aronson PL, Thurm C, Alpern ER, et al; Febrile Young Infant Research Collaborative. Variation in care of the febrile young infant <90 days in US pediatric emergency departments. *Pediatrics*. 2014;134(4):667–677
- Aronson PL, Thurm C, Williams DJ, et al; Febrile Young Infant Research Collaborative. Association of clinical practice guidelines with emergency department management of febrile infants ≤56 days of age. *J Hosp Med*. 2015;10(6):358–365
- Chua KP, Neuman MI, McWilliams JM, Aronson PL; Febrile Young Infant Research Collaborative. Association between clinical outcomes and hospital guidelines for cerebrospinal fluid testing in febrile infants aged 29-56 days. *J Pediatr*. 2015;167(6):1340–1346.e9
- Jaskiewicz JA, McCarthy CA, Richardson AC, et al. Febrile infants at low risk for serious bacterial infection—an appraisal of the Rochester criteria and implications for management. Febrile Infant Collaborative Study Group. *Pediatrics*. 1994;94(3):390–396
- Scarfone R, Murray A, Gala P, Balamuth F. Lumbar puncture for all febrile infants 29-56 days old: a retrospective cohort reassessment study. *J Pediatr*. 2017;187:200–205.e1
- de Louvois J, Halket S, Harvey D. Neonatal meningitis in England and Wales: sequelae at 5 years of age. *Eur J Pediatr*. 2005;164(12):730–734
- Libster R, Edwards KM, Levent F, et al. Long-term outcomes of group B streptococcal meningitis. *Pediatrics*. 2012;130(1). Available at: www.pediatrics.org/cgi/content/full/130/1/e8
- Pingree EW, Kimia AA, Nigrovic LE. The effect of traumatic lumbar puncture on hospitalization rate for febrile infants 28 to 60 days of age. *Acad Emerg Med*. 2015;22(2):240–243
- Lyons TW, Cruz AT, Freedman SB, et al; Pediatric Emergency Medicine Clinical Research Network (PEM CRC) Herpes Simplex Virus Study Group. Interpretation of cerebrospinal fluid white blood cell counts in young infants with a traumatic lumbar puncture. *Ann Emerg Med*. 2017;69(5):622–631
- Nigrovic LE, Kuppermann N, Neuman MI. Risk factors for traumatic or unsuccessful lumbar punctures in children. *Ann Emerg Med*. 2007;49(6):762–771
- Leazer R, Erickson N, Paulson J, et al. Epidemiology of cerebrospinal fluid cultures and time to detection in term infants. *Pediatrics*. 2017;139(5):e20163268
- Paxton RD, Byington CL. An examination of the unintended consequences of the rule-out sepsis evaluation: a parental perspective. *Clin Pediatr (Phila)*. 2001;40(2):71–77
- DeAngelis C, Joffe A, Wilson M, Willis E. Iatrogenic risks and financial costs of hospitalizing febrile infants. *Am J Dis Child*. 1983;137(12):1146–1149
- Kozyrskij AL, Ernst P, Becker AB. Increased risk of childhood asthma from antibiotic use in early life. *Chest*. 2007;131(6):1753–1759
- Trasande L, Blustein J, Liu M, Corwin E, Cox LM, Blaser MJ. Infant antibiotic exposures and early-life body mass. *Int J Obes (Lond)*. 2013;37(1):16–23
- Bailey LC, Forrest CB, Zhang P, Richards TM, Livshits A, DeRusso PA. Association of antibiotics in infancy with early childhood obesity. *JAMA Pediatr*. 2014;168(11):1063–1069
- Aronson PL, Fraenkel L. Is shared decision-making the right approach for febrile infants? *Pediatrics*. 2017;140(3):e20170225
- Byington CL, Reynolds CC, Korgenski K, et al. Costs and infant outcomes after implementation of a care process model for febrile infants. *Pediatrics*. 2012;130(1). Available at: www.pediatrics.org/cgi/content/full/130/1/e16

To Spinal Tap or Not To Spinal Tap, That Is the Question

Thomas J. Lee and Paul L. Aronson

Hospital Pediatrics 2018;8;236

DOI: 10.1542/hpeds.2017-0207 originally published online March 22, 2018;

Updated Information & Services

including high resolution figures, can be found at:
<http://hosppeds.aappublications.org/content/8/4/236>

References

This article cites 21 articles, 7 of which you can access for free at:
<http://hosppeds.aappublications.org/content/8/4/236#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Emergency Medicine
http://classic.hosppeds.aappublications.org/cgi/collection/emergency_medicine_sub
Hospital Medicine
http://classic.hosppeds.aappublications.org/cgi/collection/hospital_medicine_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://classic.hosppeds.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://classic.hosppeds.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™





To Spinal Tap or Not To Spinal Tap, That Is the Question

Thomas J. Lee and Paul L. Aronson

Hospital Pediatrics 2018;8;236

DOI: 10.1542/hpeds.2017-0207 originally published online March 22, 2018;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://hosppeds.aappublications.org/content/8/4/236>

Hospital Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 2012. Hospital Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2018 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 2154-1663.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

