

BRIEF REPORT

COVID-19: A UK Children's Hospital Experience

Hari Krishnan Kanthimathinathan, MD,^{ab} Amrit Dhesi, MBChB, MRCPCH,^a Stuart Hartshorn, FRCPCH,^{ab} Syed Habib Ali, FRCPCH,^a Jeremy Kirk, MD, FRCP, FRCPCH, DGH,^{ac} Prasad Nagakumar, MD,^{ab} Deepthi Jyothish, FRCPCH^a

OBJECTIVES: Reports from China relating to coronavirus disease (COVID-19) in children indicate a milder disease course compared with adults. Although a few pediatric COVID-19 reports from other parts of the world exist, there are none from the United Kingdom. We describe the clinical characteristics of children with COVID-19 admitted to a specialist children's hospital in United Kingdom.

METHODS: Retrospective case-series of inpatients with a positive polymerase chain reaction test for severe acute respiratory syndrome coronavirus 2, during a 6-week period from March 14 to April 24, 2020.

RESULTS: Forty-five children tested positive for severe acute respiratory syndrome coronavirus 2 during the study period. Median (interquartile range) age was 3.5 (0.7–12) years, and 31 (69%) were male. Children with comorbidities constituted 64% (29 of 45) of the study population, including 44% (20 of 45) who were considered “extremely vulnerable.” Fever (67%) and cough (55%) were the most common symptoms. High C-reactive protein (>10 mg/L) was observed in 68% (19 of 28). Lymphopenia (<1.2 × 10⁹/L) was observed in 23% (9 of 40) of children, but it was related to coexisting medical conditions in 6 children. Nine children required supplemental oxygen, two of whom received high-flow nasal cannula oxygen; one needed noninvasive ventilation and one child required invasive mechanical ventilation. Median length of stay of children with an admission outcome (*n* = 42, 93%) was 3 (2–7) days. There were no COVID-19-related deaths.

CONCLUSIONS: COVID-19 had a relatively mild course of illness in majority of the hospitalized children that included a subgroup of vulnerable children with significant comorbidities. Confirmation of this in larger nationwide studies of children is required.

ABSTRACT

www.hospitalpediatrics.org

DOI:https://doi.org/10.1542/hpeds.2020-000208

Copyright © 2020 by the American Academy of Pediatrics

Address correspondence to Deepthi Jyothish, FRCPCH, Department of General Paediatrics, Birmingham Children's Hospital, Steelhouse Lane, Birmingham B4 6NH, United Kingdom.

HOSPITAL PEDIATRICS (ISSN Numbers: Print, 2154-1663; Online, 2154-1671).

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

Drs Nagakumar, Jyothish, Kanthimathinathan, and Hartshorn and Prof Kirk conceptualized and designed the study, critically reviewed the manuscript, and approved the final manuscript as submitted; Dr Kanthimathinathan wrote the first draft of the manuscript and critically reviewed the manuscript for important intellectual content; Dr Nagakumar wrote the data analysis plan and helped with data analyses and critically reviewed the manuscript for important intellectual content; Drs Dhesi, Ali, and Hartshorn performed data collection, performed initial analyses, reviewed and revised the manuscript, and approved the final manuscript as submitted; Dr Jyothish designed the data collection instruments, coordinated and supervised data collection, and critically reviewed the manuscript for important intellectual content; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Deidentified individual participant data will not be made available.

^aBirmingham Children's Hospital Women's and Children's National Health Services Foundation Trust, Birmingham, United Kingdom; ^bBirmingham Clinical Trials Unit, University of Birmingham, Birmingham, United Kingdom; and ^cNational Institute for Health Research, Clinical Research, West Midlands, United Kingdom

Coronavirus disease 2019 (COVID-19) due to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus was declared a pandemic in March 2020.¹ Since then, COVID-19 has affected >3 million people worldwide and has been associated with >200 000 deaths worldwide.² Recent COVID-19 literature from China, including a pediatric systematic review, revealed that many children experienced a milder course of illness and better prognosis than adults.^{3,4} The review highlighted the lack of clinical or epidemiological data in children from Europe and America. There have been a few subsequent publications but no pediatric data from the United Kingdom,⁵⁻⁸ one of the worst affected countries worldwide with >25 000 fatalities.²

We describe clinical characteristics of children with COVID-19 admitted to a large regional specialist children's hospital in the West Midlands, United Kingdom. The hospital provides tertiary and quaternary specialist pediatric services, with an emergency department, with ≥ 300 pediatric inpatient beds, including a 31-bedded PICU.

METHODS

We performed a retrospective review of hospital inpatients with a positive test for SARS-CoV-2 during a 6-week period (March 14th–April 24th, 2020). Data related to demographic, clinical, laboratory, and radiologic features were obtained from hospital databases. Institutional research governance approval was obtained for waiver of ethics application, as an observational study using routinely collected data. Criteria for performing SARS-CoV-2 testing evolved during the study period but predominantly required a febrile respiratory illness (ie, temperature $\geq 37.8^{\circ}\text{C}$ and cough, coryza, shortness of breath, hoarseness, wheeze, sneezing, or respiratory distress). Testing criteria subsequently included atypical symptoms (acute abdominal pain, diarrhea, respiratory symptoms or signs without fever), especially in those requiring PICU admission, where the history was unclear or unavailable (eg admission after major trauma or burns), and immunocompromised children. Children

with a previous negative test for SARS-CoV-2 were retested if there was clinical deterioration, unexplained by an alternative diagnosis. In our study, we included children admitted from the community via the emergency department, those transferred from other hospitals, and inpatients admitted for other indications who subsequently developed symptoms meeting the above testing criteria. SARS-CoV-2 testing was performed by using real-time reverse transcriptase polymerase chain reaction from an upper airway sample (eg, combined nose and throat swabs) but occasionally from the lower respiratory tract via tracheal aspiration, if intubated. Other investigations and clinical management were performed as per relevant clinicians' discretion. Those children categorized as "extremely vulnerable" met the Public Health England UK criteria.⁹ Briefly, this includes children with various malignancies receiving chemotherapy, immunotherapy, or radiotherapy; children with recent stem cell transplant; children with respiratory conditions (eg, cystic fibrosis, severe asthma); children with rare diseases including inborn errors of metabolism; and children receiving immunosuppressive medications.⁹ Data about other underlying chronic medical conditions (eg, cerebral palsy, Down syndrome) were also recorded. Data analyses was performed by using Microsoft Excel (Microsoft Corporation, Redmond, WA), and results are presented as numbers (percentages) or median (interquartile ranges [IQRs]) as appropriate.

RESULTS

Forty-five of the 389 (12%) children tested were positive for SARS-CoV-2 antigen during the study period. Thirty-one (69%) were male. The median age of the children with COVID-19 was 3.5 (IQR: 0.7–12) years; 15 (33%) were infants (<1 year), and 14 (31%) were aged ≥ 10 years. Of those with known ethnicity data ($n = 39$): 41% were white, 28% were Asian, 13% were black, and 18% belonged to other ethnicities. Twenty-three (50%) children tested positive for SARS-CoV-2 during the third week of the study period. Clinical features of the cohort are described

in Table 1. All 3 children presenting with seizures had a previous seizure history (1 with confirmed epilepsy) and recovered rapidly with no residual neurologic deficits. A total of 64% (29 of 45) of the children in this study had at least one preexisting chronic medical condition, with 44% (20 of 45) meeting the "extremely vulnerable" Public Health England definition.

High C-reactive protein (CRP; >10 mg/L) in 68% (19 of 28) and lymphopenia ($<1.2 \times 10^9/\text{L}$) in 23% (9 of 40) were found in those children having the test performed. Of these, 8 children with elevated CRP had either coinfections, sickle cell crisis, or a recent operation or procedure, whereas 6 children with lymphopenia had malignancy-related bone marrow suppression. When children with other causes mentioned above were removed, a high CRP was observed in 55% (11 of 20) and lymphopenia was observed in 9% (3 of 34).

None of the 45 children had features of multisystem hyperinflammation or Kawasaki disease, hypotension, or shock. No specific COVID-19 drug treatments, such as immunomodulators or antiviral agents, were used in any of the study population. Three children (including 2 infants), 2 of whom had comorbidities, required high-flow nasal cannula oxygen ($n = 2$) or noninvasive continuous positive airway pressure (CPAP) ($n = 1$). One infant, with preexisting congenital heart disease, required PICU admission for invasive mechanical ventilation.

Forty-two of the 45 patients (93%) were discharged alive, and the 3 remaining hospital inpatients are there for treatment of underlying comorbidity rather than COVID-19. Median length of stay of children with an admission outcome was 3 (IQR: 2–7) days. There were no COVID-19–related deaths.

DISCUSSION

With our data, we describe a relatively mild course of COVID-19 in the vast majority of pediatric hospital inpatients. Only 20% required supplemental oxygen, and only 2 children needed any form of mechanical ventilation. Our study cohort included 44% of children within the high-risk category,

TABLE 1 Clinical Characteristics of COVID-19 in Children

Clinical Characteristics	No. (%) or Median (IQR)
Symptoms at presentation, ^a <i>n</i> (%)	
Fever	30 (67)
Cough	25 (55)
Shortness of breath	15 (33)
Gastrointestinal symptoms	13 (29)
Seizure	3 (7)
Comorbidities, ^a <i>n</i> (%)	
Hematology or oncology	10 (22)
Chromosomal or genetic disorder	9 (20)
Respiratory	6 (13)
Cardiac	4 (9)
Neurologic	4 (9)
Renal	3 (7)
Inherited metabolic disorders	1 (2)
None	16 (36)
Investigations, median (IQR)	
CRP (<i>n</i> = 28), mg/L	41.5 (2–70)
CRP (<i>n</i> = 20, excluding children with coexisting conditions), mg/L	15.5 (1–50)
Lymphocytes (<i>n</i> = 40), × 10 ⁹ /L	2.2 (1.2–4.1)
Lymphocytes (<i>n</i> = 34, excluding children with coexisting conditions), × 10 ⁹ /L	2.5 (1.6–4.8)
Chest radiograph findings (<i>n</i> = 18), <i>n</i> (%)	
Perihilar bronchial wall thickening	5 (28)
Consolidation	3 (17)
Lobar collapse	2 (11)
Respiratory support, <i>n</i> (%)	
Any supplemental oxygen	9 (20)
High-flow oxygen therapy or CPAP	3 (7)
Invasive ventilation	1 (2)

^a Children may have had multiple symptoms and/or comorbidities.

seizures) may be unrelated to COVID-19. Only 41% were white, consistent with local authority ethnicity data.¹⁴

Lymphopenia was observed in 23%, comparable to the 3.5% to 31% reported elsewhere.^{10,11} We observed elevated CRP >10 mg/L in 68%, compared with 10% to 20% in other studies,^{10,11} and although researchers of 1 study have reported ≤35% of children with elevated CRP, this used a lower CRP cutoff (>3 mg/L).¹³ Some of our study population had preexisting chronic medical conditions that potentially invalidate direct comparisons of these blood tests. After children with coexisting conditions were excluded, median CRP was only 15 (IQR: 1–50) mg/L. Moreover, only 40 of 45 had lymphocytes measured and 28 of 45 children had a CRP measurement, which may also reflect selection bias. Similarly, only 40% of children had a radiologic investigation, with most chest radiograph findings nonspecific and concurring with other similar reports.⁸ Median length of stay in this study was significantly lower than the reported mean of 12.9 days; however, discharge criteria may not be comparable, and the mean may have been skewed by a small number of long-stay patients.¹³

It is noteworthy that although only one child in our study population required intensive care support, during the same study period, >800 adults required ICU admission in the same region, demonstrating the difference in illness severity between adults and children.¹⁵

This is a retrospective report with specific inclusion criteria of pediatric inpatients positive for SARS-CoV-2 at a single UK children's hospital. Therefore, our findings are not necessarily generalizable or representative of pediatric COVID-19 in the community or, indeed, in nonspecialist hospitals. Given the relatively narrow testing criteria, asymptomatic children and children with atypical signs and symptoms may undoubtedly have been missed. We did not undertake contact tracing or ascertain the epidemiology of acquisition of COVID-19 in our study population. Nonetheless, this report represents the early experience of the clinical spectrum of COVID-19 in a

and it is reassuring that no children within this group had severe COVID-19 illness requiring PICU admission for either ventilatory or vasopressor support. The clinical course in those with underlying chronic medical conditions was similar to those without any underlying chronic medical condition, namely a mild, self-limiting illness and a short COVID-19-related inpatient stay.

Children aged 1 month to 17 years were represented in this study, as with other large-scale epidemiological studies from China, the United States, and Italy.^{3,6,8} It is reported that younger infants may be more at risk for severe illness.³ Although this current report is limited by the small

numbers of children with severe illness requiring respiratory support, 3 of the 4 requiring either high-flow nasal cannula, CPAP, or invasive ventilation were infants <1 year. In this study, 67% of children were febrile and 55% had cough, but this may reflect the testing criteria for SARS-CoV-2 in the hospital rather than the true symptomatic prevalence among children with COVID-19. Reports from China and the United States, where testing criteria included asymptomatic contacts, show 36% to 60% of children with fever and 19% to 65% had cough.^{6,8,10–13} It is possible that some children in our series could be asymptomatic carriers of SARS-CoV-2 and that their presenting symptoms (eg

pediatric cohort including complex and high-risk patients at a stand-alone children's hospital with multiple subspecialties. This is the first UK report of its kind, and although the sample size is modest, there is only one other larger cohort in the COVID-19 systematic review in children.⁴

CONCLUSIONS

We observed a relatively mild course of illness in the majority of children with COVID-19, including in those with significant comorbidities and those considered clinically extremely vulnerable. The clinical course was unaffected, despite relatively higher numbers of patients with an elevated CRP level. Most patients had prompt improvement in clinical signs and symptoms as reflected by the relatively short hospital inpatient stay. Further large-scale nationwide clinical characterization studies on COVID-19 are underway and will hopefully shed more light on this emerging illness.

Acknowledgments

We thank Drs Mark Velangi, Jim Gray, and Mitul Patel from Birmingham Women's and Children's National Health Services Foundation Trust for their contribution. Input from colleagues in the departments of cardiology, endocrinology, inherited metabolic disease, neurology, oncology, pediatric surgery, and renal medicine and staff at research and development department at Birmingham Women's and Children's National Health Services Foundation Trust is also gratefully acknowledged.

REFERENCES

1. World Health Organization. WHO Director-General's opening remarks at the media briefing on COVID-19 – 11 March 2020. Available at: <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19--11-march-2020>. Accessed April 28, 2020
2. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis*. 2020; 20(5):533–534
3. Dong Y, Mo X, Hu Y, et al. Epidemiology of COVID-19 among children in China. *Pediatrics*. 2020;145(6):e20200702
4. Castagnoli R, Votto M, Licari A, et al. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in children and adolescents: a systematic review [published online ahead of print April 22, 2020]. *JAMA Pediatr*. doi: 10.1001/jamapediatrics.2020.1467
5. Tagarro A, Epalza C, Santos M, et al. Screening and severity of coronavirus disease 2019 (COVID-19) in children in Madrid, Spain [published online ahead of print April 8, 2020]. *JAMA Pediatr*. doi: 10.1001/jamapediatrics.2020.1346
6. CDC COVID-19 Response Team. Coronavirus disease 2019 in children - United States, February 12-April 2, 2020. *MMWR Morb Mortal Wkly Rep*. 2020; 69(14):422–426
7. Gudbjartsson DF, Helgason A, Jonsson H, et al. Spread of SARS-CoV-2 in the Icelandic population. *N Engl J Med*. 2020; 382(24):2302–2315
8. Parri N, Lenge M, Buonsenso D. Children with Covid-19 in pediatric emergency departments in Italy. *N Engl J Med*. 2020; 383(2):187–190
9. Public Health England. COVID-19: guidance on shielding and protecting people defined on medical grounds as extremely vulnerable. Available at: <https://www.gov.uk/government/publications/guidance-on-shielding-and-protecting-extremely-vulnerable-persons-from-covid-19>. Accessed April 28, 2020
10. Lu X, Zhang L, Du H, et al; Chinese Pediatric Novel Coronavirus Study Team. SARS-CoV-2 infection in children. *N Engl J Med*. 2020;382(17):1663–1665
11. Qiu H, Wu J, Hong L, Luo Y, Song Q, Chen D. Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *Lancet Infect Dis*. 2020;20(6):689–696
12. Xu Y, Li X, Zhu B, et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat Med*. 2020;26(4): 502–505
13. Xia W, Shao J, Guo Y, Peng X, Li Z, Hu D. Clinical and CT features in pediatric patients with COVID-19 infection: different points from adults. *Pediatr Pulmonol*. 2020;55(5):1169–1174
14. Office for National Statistics. Census aggregate data. 2011. Available at: <http://dx.doi.org/10.5257/census/aggregate-2011-2>. Accessed April 28, 2020
15. ICNARC. ICNARC report on COVID-19 in critical care. Available at: <https://www.icnarc.org/DataServices/Attachments/Download/c5a62b13-6486-ea11-9125-00505601089b>. Accessed April 28, 2020

COVID-19: A UK Children's Hospital Experience

Hari Krishnan Kanthimathinathan, Amrit Dhesi, Stuart Hartshorn, Syed Habib Ali,
Jeremy Kirk, Prasad Nagakumar and Deepthi Jyothish

Hospital Pediatrics 2020;10;802

DOI: 10.1542/hpeds.2020-000208 originally published online June 9, 2020;

Updated Information & Services	including high resolution figures, can be found at: http://hosppeds.aappublications.org/content/10/9/802
Supplementary Material	Supplementary material can be found at:
References	This article cites 9 articles, 1 of which you can access for free at: http://hosppeds.aappublications.org/content/10/9/802#BIBL
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.hosppeds.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.hosppeds.aappublications.org/site/misc/reprints.xhtml

Hospital Pediatrics®

AN OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

COVID-19: A UK Children's Hospital Experience

Hari Krishnan Kanthimathinathan, Amrit Dhesi, Stuart Hartshorn, Syed Habib Ali,
Jeremy Kirk, Prasad Nagakumar and Deepthi Jyothish

Hospital Pediatrics 2020;10;802

DOI: 10.1542/hpeds.2020-000208 originally published online June 9, 2020;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://hosppeds.aappublications.org/content/10/9/802>

Hospital Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Hospital Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2020 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

