

# Beyond the Nursery: Postnatal Care in the 21st Century

Valerie J. Flaherman, MD, MPH,<sup>a</sup> Anthony E. Burgos, MD, MPH,<sup>b</sup> James A. Taylor, MD,<sup>c</sup> Carrie A. Phillipi, MD, PHD<sup>d</sup>

Postnatal care has been one of the great public health successes of the past century. In 1910, most infants were born at home, and perinatal mortality was high; 1 in 10 US infants died before their first birthday.<sup>1</sup> By 1960, the large majority of US births occurred in a hospital, where infants received multiple effective preventive interventions that led to a steep decrease in mortality rates. At that time, newborn preventive interventions were typically provided in a room called the “nursery,” where infants were lined up in row after row, and nurses moved from infant to infant to deliver care; mothers were housed on the postpartum ward, where they recovered from the birth process by resting quietly in bed.

Things have changed for the better in the 21st century: the number, scope, and effectiveness of postnatal preventive interventions has continued to expand, whereas infant mortality has continued to decline. We have learned that the cohousing of infants and mothers leads to better health outcomes, and providers have therefore adopted new patterns of care.<sup>2,3</sup> The preventive services previously located in the nursery now occur in a room shared between the mother and infant, and the big nursery room has gradually emptied as mothers and infants have begun to “room in.” This change has been facilitated by advocacy from programs such as the Baby Friendly Hospital Initiative<sup>4</sup> and by mounting evidence that rooming-in with mothers improves not only breastfeeding but also infant temperature regulation and glycemic control.<sup>5-7</sup>

Difficulties with the transition to extrauterine life remain common, of course, and the evidence that these transitions are ameliorated when newborns are in close proximity to their mothers<sup>8</sup> has impelled pediatric clinical staff to deliver a higher level of acuity of care in the room shared by the mother and infant than formerly. For example, intravenous antibiotics or observation and management of narcotic abstinence syndrome may require acute inpatient management but may no longer be an indication to remove the infant from the room shared with the mother.<sup>9,10</sup> This new emphasis on maintaining rooming in for infants requiring antibiotics, phototherapy, or management of other acute illnesses demands increasing expertise from physicians and nurses caring for such infants. Newborn care providers also now need enhanced expertise to provide maternal education about important care practices such as effective breastfeeding techniques, safe sleep position, tobacco exposure avoidance, and infant safety, which require reinforcement multiple times a day when mother and infant are rooming-in.

As the level of care provided in the shared mother-infant room has increased, and the nursery room itself has emptied, the term “well newborn nursery” has stuck and is still used to describe the doctors and nurses providing

[www.hospitalpediatrics.org](http://www.hospitalpediatrics.org)

DOI:<https://doi.org/10.1542/hpeds.2017-0070>

Copyright © 2017 by the American Academy of Pediatrics

<sup>a</sup>Department of Pediatrics, University of California San Francisco, San Francisco, California;

<sup>b</sup>Department of Pediatrics, Kaiser Permanente, Downey, California; <sup>c</sup>Division of General Pediatrics, School of Medicine, University of Washington, Seattle, Washington; and

<sup>d</sup>Division of General Pediatrics, Oregon Health and Science University, Portland, Oregon

Address correspondence to Valerie J. Flaherman, Department of Pediatrics, University of California San Francisco, 3333 California St, Suite 245, San Francisco, CA 94118. E-mail: [valerie.flaherman@ucsf.edu](mailto:valerie.flaherman@ucsf.edu)

HOSPITAL PEDIATRICS (ISSN Numbers: Print, 2154-1663; Online, 2154-1671).

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** No external funding.

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

Dr Flaherman conceived this commentary and wrote the first draft; Drs Burgos, Taylor, and Phillipi revised the commentary for important content; and all authors approved the final manuscript as submitted.

Opinions expressed in these commentaries are those of the authors and not necessarily those of the American Academy of Pediatrics or its Committees.

postnatal care; the room shared by the infant and mother is still often described as the postpartum room. Use of this antiquated terminology seems to lead to 2 areas of misunderstanding. First, the empty nursery room may suggest that modern medicine has made the care previously provided in the nursery obsolete. Such an observation would of course be erroneous; routine postnatal care has indeed shifted in location but has grown in importance. Second, the empty nursery room may suggest that the newborns rooming-in with mothers are well, whereas in fact, the medical staff is often exerting extensive effort to maintain a newborn's physical proximity to his or her mother when inpatient treatments such as intravenous therapy are required. For these reasons, the time has come to revise the nomenclature of this field.

We propose a sunseting of the terms nursery and well newborn unit, because they no longer sufficiently describe the location of or the care provided to newborns who are rooming-in. As an alternative, we would like to propose the terms "mother-infant unit" or "postnatal unit" to describe the location of newborns who do not require intensive care. We additionally propose the term "newborn medicine provider" to describe those who provide routine postnatal care to newborns and the term "procedure room" to describe the location where infants stay separately from their mothers when continuous observation is required. We believe such nomenclature provides a more accurate description of current postnatal care, in

keeping with the responsibility of newborn medicine providers to improve the evidence base for cost-effective care for the vast majority of US births.

Over the coming years, it is highly likely we will see even greater advances in the routine postnatal care of newborns. Retaining the term nursery risks the false connotation of antiquated images, even of an empty room, and may denigrate an important area of hospital medicine. In contrast, using terminology parallel with obstetrics, such as postnatal unit and mother-infant room, may help clarify the role of the clinical staff and of the location formerly called the nursery, and it may underscore the fundamental importance of postnatal care for promoting population health.

## REFERENCES

1. Matthews TJ, MacDorman MF, Thoma ME. Infant mortality statistics from the 2013 period linked birth/infant death data set. *Natl Vital Stat Rep*. 2015;64(9):1–30
2. Brodribb W, Kruske S, Miller YD. Baby-friendly hospital accreditation, in-hospital care practices, and breastfeeding. *Pediatrics*. 2013;131(4):685–692
3. Holmes AV, Atwood EC, Whalen B, et al. Rooming-in to treat neonatal abstinence syndrome: improved family-centered care at lower cost. *Pediatrics*. 2016; 137(6):e20152929
4. United Nations Children's Fund, World Health Organization, Wellstart International. *Baby-Friendly Hospital Initiative: Revised, Updated and Expanded for Integrated Care, Section 1, Background and Implementation, Preliminary Version*. Geneva, Switzerland: WHO Document Production Services; 2006
5. Moore ER, Anderson GC. Randomized controlled trial of very early mother-infant skin-to-skin contact and breastfeeding status. *J Midwifery Womens Health*. 2007;52(2):116–125
6. Nimbalkar SM, Patel VK, Patel DV, Nimbalkar AS, Sethi A, Phatak A. Effect of early skin-to-skin contact following normal delivery on incidence of hypothermia in neonates more than 1800 g: randomized control trial. *J Perinatol*. 2014;34(5):364–368
7. Chi Luong K, Long Nguyen T, Huynh Thi DH, Carrara HP, Bergman NJ. Newly born low birthweight infants stabilise better in skin-to-skin contact than when separated from their mothers: a randomised controlled trial. *Acta Paediatr*. 2016;105(4):381–390
8. Moore ER, Bergman N, Anderson GC, Medley N. Early skin-to-skin contact for mothers and their healthy newborn infants. *Cochrane Database Syst Rev*. 2016;11:CD003519
9. Mukhopadhyay S, Taylor JA, Von Kohorn I, et al. Variation in sepsis evaluation across a national network of nurseries. *Pediatrics*. 2017;139(3):e20162845
10. Bogen DL, Whalen BL, Kair LR, Vining M, King BA. Wide variation found in care of opioid-exposed newborns. *Acad Pediatr*. 2017;17(4):374–380

## Beyond the Nursery: Postnatal Care in the 21st Century

Valerie J. Flaherman, Anthony E. Burgos, James A. Taylor and Carrie A. Phillipi

*Hospital Pediatrics* 2017;7;763

DOI: 10.1542/hpeds.2017-0070 originally published online November 17, 2017;

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://hosppeds.aappublications.org/content/7/12/763">http://hosppeds.aappublications.org/content/7/12/763</a>
<b>Supplementary Material</b>	Supplementary material can be found at:
<b>References</b>	This article cites 9 articles, 3 of which you can access for free at: <a href="http://hosppeds.aappublications.org/content/7/12/763#BIBL">http://hosppeds.aappublications.org/content/7/12/763#BIBL</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Fetus/Newborn Infant</b> <a href="http://www.hosppeds.aappublications.org/cgi/collection/fetus:newborn_infant_sub">http://www.hosppeds.aappublications.org/cgi/collection/fetus:newborn_infant_sub</a> <b>Hospital Medicine</b> <a href="http://www.hosppeds.aappublications.org/cgi/collection/hospital_medicine_sub">http://www.hosppeds.aappublications.org/cgi/collection/hospital_medicine_sub</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.hosppeds.aappublications.org/site/misc/Permissions.xhtml">http://www.hosppeds.aappublications.org/site/misc/Permissions.xhtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://www.hosppeds.aappublications.org/site/misc/reprints.xhtml">http://www.hosppeds.aappublications.org/site/misc/reprints.xhtml</a>

# Hospital Pediatrics®

AN OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Beyond the Nursery: Postnatal Care in the 21st Century**

Valerie J. Flaherman, Anthony E. Burgos, James A. Taylor and Carrie A. Phillipi

*Hospital Pediatrics* 2017;7;763

DOI: 10.1542/hpeds.2017-0070 originally published online November 17, 2017;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://hosppeds.aappublications.org/content/7/12/763>

Hospital Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Hospital Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

