

## Supplemental Information

**SUPPLEMENTAL TABLE 3** FNASS

Systems	Signs and Symptoms	Score	AM						PM						Daily Weight
			2	4	6	8	10	12	2	4	6	8	10	12	
Central nervous system disturbances	High-pitched cry	2													
	Continuous high-pitched cry	3													
	Sleeps <1 h after feeding	3													
	Sleeps <2 h after feeding	2													
	Hyperactive Moro reflex	2													
	Markedly hyperactive Moro reflex	3													
	Mild tremors disturbed	2													
	Moderate severe tremors disturbed	3													
	Mild tremors undisturbed	1													
	Moderate severe tremors undisturbed	2													
	Increased muscle tone	2													
	Excoriation (specify area):	1													
	Myoclonic jerks	3													
	Generalized convulsions	3													
Metabolic vasomotor and respiratory disturbances	Sweating	1													
	Fever <101°F (39.3°C)	1													
	Fever >101°F (39.3°C)	2													
	Frequent yawning (>3–4 times/interval)	1													
	Mottling	1													
	Nasal stuffiness	1													
	Sneezing (>3–4 times/interval)	1													
	Nasal flaring	2													
	Respiratory rate >60/min	1													
Respiration rate >60/min with retractions	2														
Gastrointestinal disturbances	Excessive sucking	1													
	Poor feeding	2													
	Regurgitation	2													
	Projectile vomiting	3													
	Loose stools	2													
	Watery stools	3													
Summary	Total score														
	Scorer's initials														
	Status of therapy														

Adapted from Finnegan L. Neonatal abstinence syndrome: assessment and pharmacotherapy. In: Rubaltelli FF, Granti B, eds. *Neonatal Therapy: An Update*. New York, NY: Elsevier Science; 1986:122–146.